

# Instantaneous Interactions in Standard Model

A.Z. Dubničková<sup>1</sup>, S. Dubnička<sup>2</sup>, V.N. Pervushin<sup>3</sup>, M. Sečanský<sup>2,3</sup>

<sup>1</sup>Department of Theoretical Physics, Comenius University, Bratislava, Slovak Republic

<sup>2</sup>Institute of Physics, Slovak Academy of Sciences, Bratislava, Slovak Republic

<sup>3</sup>Joint Institute for Nuclear Research, 141980 Dubna, Russia

2nd February 2008

## Abstract

Instantaneous interactions in Standard Model are derived by the resolution of the Gauss constraints in inertial reference frames defined as irreducible representations of the Poincaré group. The relativistic covariant and gauge-invariant S-matrix is constructed with taking into account instantaneous interactions. Physical effects formed by the instantaneous interactions are discussed, including the enhancement in the  $K \rightarrow \pi$  transition probability and the nonleptonic and semileptonic kaon decays. The low-energy relations between these decay probability amplitudes and meson form factors are established. Exploiting these relations we estimate possible parameters of the differential  $K \rightarrow \pi e^+ e^-$  decay rate in the NA48/2 CERN experiment.

Invited talk at

The XVIII International Baldin Seminar on High Energy Physics

*Problems of Relativistic Nuclear Physics and Quantum Chromodynamics*

Joint Institute for Nuclear Research, Dubna, Russia, 25 – 30 September, 2006

# 1 Introduction

The instantaneous Coulomb interaction in QED is well known as a fixed experimental fact. Instantaneous interaction is one of the indispensable attributes of the Dirac Hamiltonian approach to QED [1, 2], where a variable without the canonical momentum is converted into a Coulomb “potential”. Remember that this Hamiltonian approach was used [3, 4] as the foundation of all heuristic methods of quantization of gauge theories, including the Faddeev-Popov (FP) method [5] and all its versions used now for description of Standard Model of elementary particles.

However, a contemporary reader could not find instantaneous interactions in the accepted version of Standard Model (SM) of elementary particles based on the FP method [6].

Why did the accepted version of SM omit instantaneous interactions?

What are the fundamental principles of foundation of instantaneous interactions?

What are physical results following from the instantaneous interactions omitted by the accepted version of SM?

Responses to these questions are the topics of the present paper.

In Section 2, we recall the Dirac Hamiltonian approach to gauge theories, where the instantaneous interactions was introduced by Dirac [1] in agreement with the priority of quantum principles and experimental facts. Section 3 is devoted to the statement of the problem of description of instantaneous interactions in SM. In Section 4, some experimental tests are discussed as evidences of instantaneous interactions. In Appendix A, S-matrix for bound states in electrodynamics is derived in terms of bilocal fields. In Appendix B, the problem of the chiral hadronization of QCD in the spirit of the non-Abelian generalization of the Dirac approach to QED is discussed.

## 2 Status of instantaneous interaction in gauge theories

### 2.1 The frame-depended variational principle

In order to explain the origin of the instantaneous interaction in SM, let us consider the Dirac approach [1] to QED given by the action

$$S_{\text{QED}} = \int d^4x \mathcal{L}_{\text{QED}}, \quad (1)$$

with Lagrangian

$$\mathcal{L}_{\text{QED}} = -\frac{1}{4} [\partial_\mu A_\nu - \partial_\nu A_\mu]^2 + \bar{\psi} [i\partial - m] \psi - A_\mu j^\mu. \quad (2)$$

where  $\partial = \partial^\mu \gamma_\mu$ ,  $A_\mu$  is a vector potential,  $\psi$  is the electron-positron field described by the Dirac bispinor,  $j_\mu = e\bar{\psi}\gamma_\mu\psi$  is the charge current, and  $e$  is the electron charge.

The action (1) is invariant with respect to gauge transformations [7, 8]

$$A_\mu^\lambda = A_\mu + \partial_\mu \lambda, \quad \psi^\lambda = e^{+ie\lambda}\psi, \quad \bar{\psi}^\lambda = e^{-ie\lambda}\bar{\psi}. \quad (3)$$

Physical solutions of the system of variational equations

$$\frac{\delta S_{\text{QED}}}{\delta A_\mu} = \partial_\nu [\partial^\nu A^\mu - \partial^\mu A^\nu] - j^\mu = 0 \quad (4)$$

are obtained in a specific *inertial frame of reference to initial data*<sup>1</sup> distinguished by a unit timelike vector  $n_\mu$  ( $n_\mu^2 = 1$ ). This vector splits gauge field  $A_\mu$  into the timelike  $A_0 = A_\mu n_\mu$  and spacelike  $A_\nu^\perp = A_\nu - n_\nu(A_\mu n_\mu)$  components<sup>2</sup>.

The time component of Eq. (4) with respect to  $A_0$  is the Gauss constraint

$$\Delta A_0 - \partial_0 \partial_k A_k = -j_0, \quad (5)$$

where  $\Delta = \partial_j^2$ ,  $j_0 = e\bar{\psi}\gamma_0\psi$  is the current. In accordance with the theory of differential equations, the field components  $A_0$  cannot be a *degree of freedom*, because the canonical momentum  $P_0 = \partial\mathcal{L}/\partial(\partial_0 A_0) = 0$  is equal to zero. An exact solution of Eq. (5)

$$A_0(x_0, x_k) = \frac{1}{\Delta}[\partial_k A_k - j_0] \equiv -\frac{1}{4\pi} \int d^3y \frac{[\partial_k A_k(x_0, y_k) - j_0(x_0, y_k)]}{|\vec{x} - \vec{y}|}. \quad (6)$$

is treated as the *Coulomb potential field* leading to the *instantaneous* interaction. The source of this *potential field* can be only an electric current  $j_0$ . The linear terms  $\partial_0 \partial_k A_k$  in this solution (6) cannot be considered as a physical source of the Coulomb potential. Therefore, Dirac [1] supposed to remove this term from the Gauss constraint (5) by gauge transformations

$$A_\mu^{(\text{rad})}(A) = A_\mu + \partial_\mu \Lambda^{(\text{rad})}(A), \quad \psi^{(\text{rad})}(\psi, A) = e^{ie\Lambda^{(\text{rad})}(A)}\psi, \quad (7)$$

where

$$\Lambda^{(\text{rad})}(A) = -\frac{1}{\Delta} \partial_k A_k \equiv \frac{1}{4\pi} \int d^3y \frac{\partial_k A_k(x_0, y_k)}{|x - y|}. \quad (8)$$

## 2.2 Radiation variables as Dirac's gauge-invariant observables

This transformation (7) determines new variables

$$A_0^{(\text{rad})}(A) = A_0 + \partial_0 \Lambda^{(\text{rad})}(A) = A_0 - \partial_0 \frac{1}{\Delta} \partial_k A_k, \quad (9)$$

$$A_l^{(\text{rad})}(A) = A_l + \partial_l \Lambda^{(\text{rad})}(A) = A_l - \partial_l \frac{1}{\Delta} \partial_k A_k, \quad (10)$$

$$\psi^{(\text{rad})}(\psi, A) = e^{ie\Lambda^{(\text{rad})}(A)}\psi. \quad (11)$$

Thus, the frame-fixing  $A_\mu = (A_0, A_k)$ , the treatment of  $A_0$  as a classical field, and the Dirac diagonalization (7) of the Gauss law (5) [1] lead to variables as the functionals that are invariant with respect to gauge transformations (3) of the initial fields  $A_\mu, \psi$

$$A_\mu^{(\text{rad})}(A + \partial\lambda) = A_\mu^{(\text{rad})}(A), \quad \psi^{(\text{rad})}(\psi e^{ie\lambda}, A + \partial\lambda) = \psi^{(\text{rad})}(\psi, A). \quad (12)$$

---

<sup>1</sup>The physical concept of a *frame of reference to initial data* is defined as a three-dimensional coordinate basis with a watch and a ruler (and other physical devices) for measurement of time and distance (velocity, mass, and other physical quantities), i.e. the *initial data* required for unambiguous resolving equations of motion (4). *Inertial* means that this coordinate basis is connected with a heavy physical body moving without influences of any external forces. Inertial frame of reference in Minkowskian space-time is associated with the unit time-axis  $n_\mu = (\frac{1}{\sqrt{1-\vec{v}^2}}, \frac{\vec{v}}{\sqrt{1-\vec{v}^2}})$ , where  $\vec{v}$  is the velocity. The frame of reference  $n_\mu^{\text{cf}} = (1, 0, 0, 0)$  with  $\vec{v} = 0$  is called the comoving frame. Transition to another *inertial frame* are fulfilled by a Lorentz transformation  $L_{\mu\nu} n_\mu^{\text{cf}} = n_\mu$ . A complete set of inertial frames  $\{n_\mu\}$  is obtained by all Lorentz transformations of a comoving frame.

<sup>2</sup>Recall that this specific reference frame was chosen by Wigner [9], in order to construct irreducible representations of the Poincaré group supposing the existence of a vacuum as a state with minimal energy (see in detail [10, 11]).

These functionals were called the “dressed fields”, or the “radiation variables” [1].

In terms of variables (9) – (11) the Gauss law (5) takes the diagonal form (i.e. it loses the linear term  $\partial_k A_k$ )

$$\Delta A_0^{(\text{rad})}(A) = -j_0^{(\text{rad})} \equiv -e\bar{\psi}^{(\text{rad})}\gamma_0\psi^{(\text{rad})} \quad (13)$$

and the spatial components become transversal identically

$$\partial_k A_k^{(\text{rad})}(A) \equiv 0. \quad (14)$$

After the substitution of a manifest resolution of the Gauss constraint (5) into the initial action (1)<sup>3</sup> this action can be expressed in terms of the gauge-invariant radiation variables (9) – (11) [1, 2]

$$\begin{aligned} S_{\text{QED}} \Big|_{\frac{\delta S_{\text{QED}}}{\delta A_0}=0} &\equiv S_{\text{QED}}^{(\text{rad})} = \\ &= \int d^4x \left[ \frac{1}{2}(\partial_\mu A_k^{(\text{rad})})^2 + \bar{\psi}^{(\text{rad})}[i\partial - m]\psi^{(\text{rad})} + A_k^{(\text{rad})}j_k^{(\text{rad})} - \frac{1}{2}j_0^{(\text{rad})}\frac{1}{\Delta}j_0^{(\text{rad})} \right]. \end{aligned} \quad (15)$$

One can see that Dirac in [1] managed to express his Hamiltonian scheme in terms of radiation variables (9) – (11) as functionals from the initial fields, and these functionals are invariant (12) with respect to gauge transformations of the initial fields (3). In other words, one can say that the Dirac radiation variables are manifestly gauge-invariant. Therefore, the fact that the current  $j_\mu^{(\text{rad})}$  cannot satisfy the conservation constraint

$$\partial_\mu j^{(\text{rad})\mu} \neq 0. \quad (16)$$

does not contradict the gauge-invariance of the initial action (1) and, moreover, it allows us to give the gauge-invariant description of bound states, where the current  $j_\mu^{(\text{rad})}$  is not conserved.

### 2.3 Dirac’s frame depended approach versus the frame free one

Values of the reduced action (15) onto solutions of the motion equations takes the form

$$\begin{aligned} S_{\text{QED}}^{(\text{rad})} \Big|_{\frac{\delta S_{\text{QED}}}{\delta A_k}=0} &= \frac{1}{2} \int d^4x d^4y \left[ j_0^{(\text{rad})}(x) \mathbf{C}_{00}(x-y) j_0^{(\text{rad})} + j_k^{(\text{rad})}(x) \mathbf{T}_{ki}(x-y) j_i^{(\text{rad})}(y) \right] + \\ &+ \int d^4x \bar{\psi}^{(\text{rad})}[i\partial - m]\psi^{(\text{rad})}, \end{aligned} \quad (17)$$

where

$$\mathbf{C}_{00}(x-y) = -\delta(x_0 - y_0) \frac{1}{4\pi|\vec{x} - \vec{y}|}, \quad (18)$$

$$\mathbf{T}_{ki}(x-y) = \frac{1}{(2\pi)^4} \int d^4q \left( \delta_{ki} - \frac{q_k q_i}{\vec{q}^2} \right) \frac{e^{iq_\nu(x-y)^\nu}}{q_\mu^2 + i\epsilon}. \quad (19)$$

The action (17) allows one to determine the propagator of radiation variables in the momentum representation  $j(x) = (1/(2\pi)^2) \int d^4q e^{-iq_\nu x^\nu} \tilde{j}(q)$ . One can see that the current part of the action (17) takes the form

$$S^{(\text{rad})}[j] = \frac{1}{2} \int d^4q \tilde{j}^\mu(q) \tilde{j}^\nu(q) D_{\mu\nu}^{\gamma,(\text{rad})}(q), \quad (20)$$

---

<sup>3</sup>This substitution, i.e. the calculation of value of the action onto a solution of the Gauss constraint, is called the reduction procedure. This reduction allows us to eliminate nonphysical pure gauge degrees of freedom [12].

where

$$D_{\mu\nu}^{\gamma,(\text{rad})}(q) = \delta_{\mu 0}\delta_{\nu 0}\frac{1}{\vec{q}^2} + \delta_{\mu i}\delta_{\nu j}\left(\delta_{ij} - \frac{q_i q_j}{\vec{q}^2}\right)\frac{1}{q_\mu^2 + i\varepsilon} \equiv \mathbf{C}_{\mu\nu} + \mathbf{T}_{\mu\nu} \quad (21)$$

is the propagator as the sum of the instantaneous exchange  $\mathbf{C}_{\mu\nu}$  and transverse field one  $\mathbf{T}_{\mu\nu}$ .

This propagator contains two singularities: the Coulomb one  $\mathbf{C}$  forming instantaneous atoms (see Appendix A) and the light cone ones of the transverse variables  $\mathbf{T}$  (considered as radiation corrections). This propagator can be identically rewritten as the sum of the Feynman propagator  $\mathbf{F}$  (that does not depend on the frame time-axis  $n_\mu$ ) and the longitudinal  $\mathbf{L}$  one:

$$D_{\mu\nu}^{\gamma,(\text{rad})}(q) \equiv -\frac{\delta_{\mu\nu}}{q_\nu^2 + i\varepsilon} + \frac{(q_0\delta_{\mu 0})(q_0\delta_{\nu 0}) - (q_i\delta_{\mu i})(q_j\delta_{\nu j})}{\vec{q}^2[q_\nu^2 + i\varepsilon]} \equiv \mathbf{F}_{\mu\nu}(q) + \mathbf{L}_{\mu\nu}(q). \quad (22)$$

It is easy to see that if the currents  $\tilde{j}_\mu(q)$  satisfy the conservation constraint

$$q^\mu \tilde{j}_\mu^{\text{(rad)}} = 0 \quad (23)$$

in contrast to (16), the action (20) takes the form

$$S^{(\text{rad})}[j] = \frac{1}{2} \int d^4q \tilde{j}^\mu(q) \mathbf{F}_{\mu\nu}(q) \tilde{j}^\nu(q) \equiv -\frac{1}{2} \int d^4q \frac{\tilde{j}_\mu^2(q)}{q_\nu^2 + i\varepsilon}. \quad (24)$$

This result corresponds to the Lagrangian

$$\mathcal{L}_{\text{QED}}^{\text{F}} = \mathcal{L}_{\text{QED}} - \frac{1}{2}(\partial_\mu A^\mu)^2, \quad (25)$$

where the first term is the initial Lagrangian (2). The change

$$\mathcal{L}_{\text{QED}}|_{(\partial_\mu j^\mu=0)} \Rightarrow \mathcal{L}_{\text{QED}}^{\text{F}} = \mathcal{L}_{\text{QED}} - \frac{1}{2}(\partial_\mu A^\mu)^2 \quad (26)$$

violates gauge-invariance (in contrast to the radiation variables) and it is known as the *Feynman gauge*. A similar change (26) is the basis of the accepted frame free formulation of SM [6].

Thus, in the Dirac Hamiltonian approach to QED we have perturbation theory featuring two singularities in the photon propagators (21) (the instantaneous singularity and that at the light cone), whereas the perturbation theory in the frame free gauge (26) (accepted now to formulate SM [6]) involves only one singularity in the propagators (that at the light cone (24)). The perturbation theory in this gauge in terms of the propagators with the light cone singularity can by no means describe instantaneous Coulomb atoms, and it contains only the frame free Wick-Cutkosky bound states whose spectrum is not observed in nature [13, 14].

The problem of different physical results in two different *gauges* cannot be explained by the violation of the gauge-invariance in the Lagrangian (26). We can keep gauge-invariance of the frame free method choosing the Lorentz variables  $A_\mu^{(\text{L})}(A)$  by gauge transformations

$$A_\mu^{(\text{L})}(A) = A_\mu + \partial_\mu \Lambda^{(\text{L})}(A), \quad \psi^{(\text{L})}(\psi, A) = e^{ie\Lambda^{(\text{L})}(A)}\psi, \quad (\Lambda^{(\text{L})}(A) = -\frac{1}{\square}\partial_\mu A^\mu). \quad (27)$$

The gauge-invariant functional  $A_\mu^{(\text{L})}(A)$  identically satisfies the Lorentz constraint  $\partial_\mu A^{(\text{L})\mu} \equiv 0$ .

In fact, the change from the radiation variable (7)  $A^{(\text{rad})}, \psi^{(\text{rad})}$  to the frame free Lorentz ones (27)  $A^{(\text{L})}, \psi^{(\text{L})}$

$$A_\mu^{(\text{rad})}(A^{(\text{L})}) = A_\mu^{(\text{L})} + \partial_\mu \Lambda^{(\text{rad})}(A^{(\text{L})}), \quad (28)$$

$$\psi^{(\text{rad})}(\psi^{(\text{L})}, A^{(\text{L})}) = e^{ie\Lambda^{(\text{rad})}(A^{(\text{L})})} \psi^{(\text{L})}, \quad (29)$$

where  $\Lambda^{(\text{rad})}(A^{(\text{L})})$  is the Dirac phase given by Eq. (8), keeps all physical results, if there remains the Dirac *gauge of physical sources* in terms of the new variables:

$$\mathcal{L}_{\text{sources}}^{(\text{rad})} = J_k^{(\text{rad})} [A_k^{(\text{L})} + \partial_k \Lambda^{(\text{rad})}(A^{(\text{L})})] + \bar{\eta}^{(\text{rad})} e^{ie\Lambda^{(\text{rad})}(A^{(\text{L})})} \psi^{(\text{L})} + \bar{\psi}^{(\text{L})} e^{-ie\Lambda^{(\text{rad})}(A^{(\text{L})})} \eta^{(\text{rad})}. \quad (30)$$

As a relic of the Dirac fundamental quantization, the Dirac phase factors  $e^{ie\Lambda^{(\text{rad})}(A^{(\text{L})})}$  in the source Lagrangian (30) “remember” the entire body of information about the *frame of reference to initial data* and the instantaneous interaction. As was predicted by Schwinger [16], all these effects disappear, leaving no trace, if these Dirac factors are removed by means of the substitution

$$\mathcal{L}_{\text{sources}}^{(\text{rad})} \rightarrow \mathcal{L}_{\text{sources}}^{(\text{L})} = J_\mu^{(\text{L})} A_\mu^{(\text{L})} + \bar{\eta}^{(\text{L})} \psi^{(\text{L})} + \bar{\psi}^{(\text{L})} \eta^{(\text{L})}. \quad (31)$$

This substitution (treated as another choice of *gauge of physical sources* by the elimination of the Dirac phase factors) loses the instantaneous interaction and instantaneous bound states. We see that there is the strong dependence of physical results on the *gauge of physical sources*, and this dependence does not mean the violation of the gauge-invariance principle, because both the radiation variables (7) and the Lorentz ones (27) are gauge-invariant functionals of the initial fields<sup>4</sup>.

The region of validity of the theorem equivalence of the frame dependent method and the frame free *Lorentz gauge formulations* (changing the *gauge of physical sources*) is restricted by the elementary particle scattering amplitudes [3].

The change of the *gauge of physical sources* (31) is just the point where the frame free FP method in the *Lorentz gauge* (26) loses the instantaneous interaction (18) forming instantaneous bound states (considered in Appendixes A and B). Moreover, the field  $A_0$  treated as a dynamic one in the *the Lorentz gauge formulation* (26) gives a negative contribution to energy and it leads to tachyon in the spectrum of the Wick-Cutkosky bound states [13, 14].

The standard frame free approach [6] to SM extending the FP method out of the region of its validity (23) finally lost instantaneous interactions. However, if we keep instantaneous interactions, the question arises about a choice of the frame and relativistic invariance. What do relativistic covariant instantaneous atoms in QED mean?

## 2.4 Relativistic invariance: “frame free” versus “many frames”

It is interesting to trace the historical evolution of the physical concepts of *relativistic invariance* and *gauge invariance*. Both Julian Schwinger (who “*rejected all Lorentz gauge formulations as unsuited to the role of providing the fundamental operator quantization*” [16]) and Faddeev (who accepted these *formulations* in [3]) used the same argument of *relativistic invariance*. The phrase “*Finally, the Hamiltonian formulation, in the case of field theory, is not relativistically*

---

<sup>4</sup>Therefore, the assertion that “*the dependence of physical result on the choice of a gauge means the violation of the gauge-invariance principle*” is not correct. The correct full name of the concept “gauge” is *gauge of physical sources*. A change of *gauge of physical sources* can lead to a change of physical results as we have seen above.

*covariant*' was treated in [3] as a defect of the frame dependent Hamiltonian formulation and one of the arguments in favor of the Lorentz frame free gauge (without the instantaneous interaction). However, in accord with the theory of representations of the Poincaré group the relativistic invariance means that *a complete set of states*  $\{|\Phi_I\rangle\}_{n^{\text{cf}}}$  *obtained by all Lorentz transformations of a state*  $|\Phi\rangle_0$  *in a definite inertial frame of reference*  $n_\mu^{\text{cf}} = (1, 0, 0, 0)$  *coincides with a complete set of states*  $\{|\Phi_I\rangle\}_n$  *obtained by all Lorentz transformations of this state*  $|\Phi\rangle_0$  *in another frame of reference*  $n_\mu = (\frac{1}{\sqrt{1-\vec{v}^2}}, \frac{\vec{v}}{\sqrt{1-\vec{v}^2}})$  [10]. Therefore, the frame dependence of the Hamiltonian formulation was treated in [16] as necessity to prove that the algebra of commutation relations of gauge-invariant observables coincide with the Poincaré algebra. This proof was fulfilled by Zumino [15] (and Schwinger [16] in the case of non-Abelian fields even in 1962) at the level of the algebra of commutation relations of the Hamiltonian and momentum operators with the nonlocal commutation relation  $i[A_k^{(\text{rad})}(x), P_j^{(\text{rad})}(y)] = [\delta_{kj} - \frac{\partial_k \partial_j}{\Delta}] \delta^3(x-y)$  for quantum radiation variables taken to be symmetrically ordering in the Hamiltonian.

We can see that the *relativistic invariance* as the theory of irreducible representation of the Poincaré group does not mean the frame free formulation (27). The *relativistic invariance* means that a complete set of all physical states includes the states obtained by *all Lorentz transformations of the comoving frame*, i.e. all Lorentz transformations of the instantaneous interactions obtained in the comoving frame. Thus, there are two following possibilities to construct the weak S-matrix amplitude transitions between physical states, i) the frame free S-matrix, and ii) the many-frame S-matrix.

## 2.5 Many-frame relativistic S-matrix with instantaneous interactions

Recall that S-matrix in QED is constructed on the basis of the Schrödinger equation

$$H^{(\text{rad})}|\Phi(x^0)\rangle = i\frac{d}{dx^0}|\Phi(x^0)\rangle, \quad (32)$$

where

$$H^{(\text{rad})} = \int_{V_0} d^3x \mathcal{H}^{(\text{rad})}[n] = \int_{V_0} d^3x [\mathcal{H}_0(n) + \mathcal{H}_{\text{ins.int.}}(n) + \mathcal{H}_{\text{ret.int.}}(n)], \quad (33)$$

$$\mathcal{H}^{(\text{rad})}(n) = [P_i^{(\text{rad})}\partial_0 A_i^{(\text{rad})} + P_\psi^{(\text{rad})}\partial_0 \psi^{(\text{rad})} - \mathcal{L}^{(\text{rad})}] \quad (34)$$

are the Hamiltonian and its density as a sum of free field part, instantaneous interaction part, and retarded interaction one;  $\Phi$  is a physical states. All these notions (space-time, volume, energy, class of functions and etc.) make sense only in a definite *frame of reference to initial data* distinguished by a unit time-axis  $n_\mu = (\frac{1}{\sqrt{1-\vec{v}^2}}, \frac{\vec{v}}{\sqrt{1-\vec{v}^2}})$ , where  $\vec{v}$  is the initial velocity of an inertial frame,  $\vec{v} = 0$  is called a *comoving frame* (cf) (see the footnote 1). The resolution of the equation takes the form of the operator of evolution

$$|\Phi(x^0)\rangle = T \exp \left\{ -i \int_{x_{(\text{in})}^0}^{x^0} dx^0 H^{(\text{rad})} \right\} |\Phi_I(x_{(\text{in})}^0)\rangle, \quad (35)$$

here  $|\Phi_I(x_{(\text{in})}^0)\rangle$  is an initial state defined in the asymptotic region, where the *Hamiltonian of retarded interaction* in the sum (33) can be neglected<sup>5</sup>, so that one can determine the energy

<sup>5</sup>We can neglect the retarded interaction provided that  $|x_{(\text{out})}^0 - x_{(\text{in})}^0| \gg E_{I,\text{min}}^{-1}$ ,  $V_0^{1/3} \gg E_{I,\text{min}}^{-1}$ . This condition means that all stationary solutions with zero energy  $E_{I,\text{min}} \rightarrow 0$  cannot be considered as perturbational.

spectrum  $E_I$  from the equation

$$\int d^3x [\mathcal{H}_0(n) + \mathcal{H}_{\text{ins.int.}}(n)] |\Phi_I(x_{\text{(in)}}^0) \rangle = E_I |\Phi_I(x_{\text{(in)}}^0) \rangle. \quad (36)$$

In accord with the *spectrality principle* a complete set of eigenvalues  $E_I$  (with a finite energy density  $E_I/V_0 \leq \infty$ ) of a manifold of all asymptotic states with the quantum numbers  $I$  includes a physical vacuum as a state with a minimal energy  $E_{I,\text{min}}$  [11].

Notice that in the comoving frame  $n_\mu^{\text{cf}} = (1, 0, 0, 0)$  the set of one-particle and two-particle equations<sup>6</sup> (36) coincide with the Schwinger - Dyson equation and the Salpeter one, respectively (see Appendix A).

The next step is the construction of the S-matrix elements as weak probability amplitude transitions between the all asymptotic<sup>7</sup> states including bound states. A complete set of these states includes any bound state moving with the momentum  $\mathcal{P}_\mu$  obtained the Lorentz transformation of the time-axis

$$n_\mu^{\text{cf}} = (1, 0, 0, 0) \implies n_\mu^{\mathcal{P}} = \frac{\mathcal{P}_\mu}{\sqrt{\mathcal{P}^2}} = \left( \frac{\mathcal{P}_0}{\sqrt{\mathcal{P}^2}}, \frac{\mathcal{P}_k}{\sqrt{\mathcal{P}^2}} \right). \quad (37)$$

Therefore, practically, in QED, one uses the many-frame S-matrix

$$S_{J_{\text{out}}, I_{\text{in}}}^{(\text{rad, practical})} = \left\langle \Phi_J(x_{\text{(out)}}^0) \left| \exp \left\{ -i \int d^4x \mathcal{H}^{(\text{rad})}(\hat{n}^{\text{ID}}) \right\} \right| \Phi_I(x_{\text{(in)}}^0) \right\rangle, \quad (38)$$

where the time-axis  $n_\mu^{\text{cf}} = (1, 0, 0, 0)$  [16] is replaced by *the initial datum operator*  $\hat{n}_\mu^{\text{ID}}$  proportional to the operator of the total momentum<sup>8</sup>

$$\hat{n}_\mu^{\text{ID}} |\Phi^J \rangle = \frac{P_\mu^J}{M_J} |\Phi^J \rangle \quad (39)$$

of any physical state considered as a irreducible representation of the Poincaré group [20, 21], where this operator  $\hat{n}_\mu^{\text{cf}}$  becomes c-number (see Appendix A).

Thus, instead of the acceptable frame free formulation of the Standard Model [6] without instantaneous interactions we suppose to use the Dirac-like radiation variables with instantaneous interactions and the many-frame S-matrix (38) where the comoving time-axis is replaced by the initial datum operator (39)<sup>9</sup>.

<sup>6</sup>Similar equations are well - known from the nonrelativistic many - body theory (Landau's theory of fermi liquids [17], Random Phase Approximation [18]) and play an essential role in the description of elementary excitation in atomic nuclei [19]. Their relativistic analogies describe in QCD the Goldstone pion and the constituent masses of the light quarks [20, 21, 22] (see Appendix B).

<sup>7</sup>The hypothesis of asymptotical states of S-matrix in QFT means that physical states are created together with their comoving *reference frames* where quantum numbers of these states are measured.

<sup>8</sup>It is clear that at the point of the existence of the bound state with the definite total momentum  $\mathcal{P}_{A\mu}$  any instantaneous interaction with the time-axis  $n_\mu$  parallel to this momentum is much greater than any retarded interaction in the sum (33) [23].

<sup>9</sup>In particular, this definition of S-matrix (38) justifies the application of the axial gauge for the description of the deep-inelastic scattering amplitudes, where this gauge imitates the transition to the large momentum frame [24].



### 3 Instantaneous weak interactions

#### 3.1 Standard Model

We consider the accepted SM without any additional terms violating gauge invariance

$$\mathcal{L}_\phi = \frac{1}{2}(\partial_\mu \phi)^2 - \frac{\lambda}{4}[\phi^2 - v^2]^2 - \phi[f_v(\bar{e}_R e_L + \bar{e}_L e_R) + (\bar{e}_R e_L + \bar{e}_L e_R)] + \frac{1}{8}\phi^2[g^2 W_\mu^+ W^{-\mu} + (g^2 + g'^2)Z_\mu^2] \quad (40)$$

$$\mathcal{L}_l = \bar{\nu}_e i(\gamma \partial) \nu_e + \frac{g}{2 \cos \theta_W}(\bar{\nu}_e \gamma_\mu^L \nu_e) Z^\mu + \bar{e} i \gamma^\mu (\partial_\mu + i g \sin \theta_W A_\mu) e + \frac{g}{2 \cos \theta_W}(\bar{e} \gamma_\mu^L e) Z^\mu + g \frac{\sin^2 \theta_W}{\cos \theta_W}(\bar{e} \gamma_\mu e) Z^\mu + \frac{g}{\sqrt{2}}(\bar{\nu}_e W_\mu^+ \gamma^\mu e_L + \bar{e}_L W_\mu^- \gamma^\mu \nu_e) \quad (41)$$

$$\begin{aligned} \mathcal{L}_V = & -\frac{1}{4}(\partial_\mu A_\nu - \partial_\nu A_\mu)^2 - \frac{1}{4}(\partial_\mu Z_\nu - \partial_\nu Z_\mu)^2 - \frac{1}{2}[D_\mu W_\nu^+ - D_\nu W_\mu^+]^2 \\ & - i e(\partial_\mu A_\nu - \partial_\nu A_\mu) W^{+\mu} W^{-\nu} - g^2 \cos^2 \theta_W [Z_\nu^2 (W_\mu^+ W^{-\mu}) - (W_\mu^+ Z^\mu)(W_\nu^- Z^\nu)] + \\ & + i g \cos_W \theta (\partial_\mu Z_\nu - \partial_\nu Z_\mu) W^{+\mu} W^{-\nu} + \frac{1}{2} i g \cos_W \theta [(D_\mu W_\nu^+ - D_\nu W_\mu^+)(W^{-\mu} Z^\nu - W^{-\nu} Z^\mu) - c.c.], \end{aligned} \quad (42)$$

where  $V_\mu^K = (A_\mu, Z_\mu, W_\mu^+, W_\mu^-)$  are the standard set of vector fields, and with the Higgs effect of spontaneous symmetry breaking

$$\phi = \eta + v, \quad v^2 = \frac{m_\eta^2}{2\lambda}, \quad (43)$$

where  $\eta$  is the Higgs field with the mass  $m_\eta$ .

First of all, we can see that this action (42) is bilinear with respect to the time components of the vector fields  $V_0^K = (A_0, Z_0, W_0^+, W_0^-)$  in the “comoving frame”  $n_\mu^{\text{cf}} = (1, 0, 0, 0)$

$$S_V = \int d^4x \left[ \frac{1}{2} V_0^K \hat{L}_{00}^{KI} V_0^I + V_0^K J^K + \dots \right], \quad (44)$$

where  $\hat{L}_{00}^{KI}$  is the matrix of differential operators. Therefore, the Dirac approach to SM can be realized. This means that the problems of the reduction and diagonalization of the set of the Gauss laws are solvable, and the Poincaré algebra of gauge-invariant observables can be proved.

In any case, SM in the lowest order of perturbation theory is reduced to the sum of the Abelian massive vector fields, where Dirac’s radiation variables was considered in [25].

#### 3.2 The reduction of the Abelian massive vector theory

The classical action of massive QED is

$$W = \int d^4x \mathcal{L}(x) = \int d^4x \left[ -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} M^2 V_\mu^2 + \bar{\Psi}(i\partial - m)\Psi - V_\mu J^\mu \right], \quad (45)$$

with the Lagrangian  $\mathcal{L}(x)$  and the currents  $J_\mu \equiv e\bar{\Psi}\gamma_\mu\Psi$ . The Euler-Lagrange equation for  $V_0$  is a constraint

$$\frac{\delta W}{\delta V_0} = 0 \quad \leftrightarrow \quad (\Delta - M^2)V_0 = -\partial_i \dot{V}_i + J_0, \quad (46)$$

It has the solution

$$V_0[\vec{V}, J_0] = \frac{1}{\Delta - M^2}(-\partial_i \dot{V}_i + J_0). \quad (47)$$

Inserting this into  $W$  we obtain the reduced  $W_{\text{red}}$

$$W_{\text{red}}[\vec{V}, \Psi] = \int d^4x \mathcal{L}_{\text{red}} \equiv \int d^4x (\mathcal{L}_{\text{red}}^V + \mathcal{L}_{\text{red}}^\Psi), \quad (48)$$

where

$$\begin{aligned} \mathcal{L}_{\text{red}}^V &= \frac{1}{2} \left( \dot{V}_i R_{ij} \dot{V}_j + V_i (\Delta - M^2) R_{ij} V_j \right), \\ \mathcal{L}_{\text{red}}^\Psi &= \frac{1}{2} J_0 \frac{1}{\Delta - M^2} J_0 + J_0 \left( \frac{1}{\Delta - M^2} \partial_i \dot{V}_i \right) + V_i J_i + \bar{\Psi} (i \not{\partial} - m) \Psi, \end{aligned} \quad (49)$$

are the reduced Lagrangians with the operator

$$R_{ij} \equiv \delta_{ij} - \frac{\partial_i \partial_j}{\Delta - M^2} = \delta_{ij}^T - \frac{M^2}{\Delta - M^2} \delta_{ij}^{\parallel} \quad (50)$$

where we have used the longitudinal and transverse projection operators

$$\delta_{ij}^{\parallel} \equiv \frac{\partial_i \partial_j}{\Delta}, \quad \delta_{ij}^T \equiv \delta_{ij} - \delta_{ij}^{\parallel}. \quad (51)$$

In contrast to the massless case,  $R_{ij}$  is not a projection operator,  $R^2 \neq R$ , but

$$R_{ij} R_{jl} = R_{il} + \frac{M^2 \partial_i \partial_l}{(\Delta - M^2)^2} \quad (52)$$

and  $R_{ij}$  is invertible

$$R_{ij}^{-1} = \delta_{ij}^T - \frac{\Delta - M^2}{M^2} \delta_{ij}^{\parallel} = \delta_{ij} - \frac{\partial_i \partial_j}{M^2}. \quad (53)$$

In the massless limit however the reduction operator becomes the transverse projection operator for photons and ceases to be invertible.

The second term in  $\mathcal{L}_{\text{red}}^\Psi$  (49) can be eliminated (in order to diagonalize the Gauss law) by introducing the new radiation (R) variables

$$V_j^R = \left( \delta_{ij} - \frac{\partial_i \partial_j}{\Delta - M^2} \right) V_j = R_{ij} V_j \quad (54)$$

$$\Psi^R \equiv \exp \left( i e \frac{1}{\Delta - M^2} \partial_i V_i \right) \Psi \quad (55)$$

which generalizes the Dirac radiation variables in QED (10) and (11). Since (55) is only a phase transformation, the corresponding current  $J_\mu$  stays the same and  $\mathcal{L}_{\text{red}}^\Psi$  becomes

$$\mathcal{L}_{\text{red}}^\Psi = \frac{1}{2} J_0 \frac{1}{\Delta - M^2} J_0 + J_i \left( \delta_{ij} - \frac{\partial_i \partial_j}{\Delta - M^2} \right) V_j + \bar{\Psi}^R (i \not{\partial} - m) \Psi^R. \quad (56)$$

Using the radiation variables  $V_k^R = R_{kj}V_j$  with  $R_{ij}$  given by (50) the reduced Lagrangian can be written

$$\mathcal{L}_{\text{red}} = \frac{1}{2} \left( \dot{V}_i^R R_{ij}^{-1} \dot{V}_j^R + V_i^R (\Delta - M^2) R_{ij}^{-1} V_j^R \right) + \bar{\Psi}^R (i\partial - m) \Psi^R + J_i V_i^R + \frac{1}{2} J_0 \frac{1}{\Delta - M^2} J_0. \quad (57)$$

Thus, the frame-fixing  $A_\mu = (A_0, A_k)$ , the treatment of  $A_0$  as a classical field, and the Dirac-like diagonalization of the Gauss law (46) (i.e. eliminating the linear field contribution  $\partial_i \dot{V}_i$ ) lead to radiation variables (54) and (55) that cannot be associated with any gauge constraint, in the case of the massive fields [25].

### 3.3 Dirac's frame dependent approach versus the frame free one

The Fourier transform of the reduction operator  $R_{ij}$

$$\sum_\lambda \epsilon_i^{P(\lambda)}(\vec{q}) \epsilon_j^{P(\lambda)}(\vec{q}) = R_{ik}(q) \left( \delta_{kl} + \frac{q_k q_l}{M^2} \right) R_{lj}(k) = R_{ij}(q) = \delta_{kl} - \frac{q_k q_l}{\vec{q}^2 + M^2}. \quad (58)$$

leads to the free propagator

$$\begin{aligned} D_{ij}^R(x-y) &= \langle 0 | T V_i^R(\vec{x}, x_0) V_j^R(\vec{y}, y_0) | 0 \rangle \\ &= -i \int \frac{d^4 q}{(2\pi)^4} \frac{e^{-iq \cdot (x-y)}}{q^2 - M^2 + i\epsilon} \left( \delta_{kl} - \frac{q_k q_l}{\vec{q}^2 + M^2} \right). \end{aligned} \quad (59)$$

Together with the instantaneous interaction (see the last term in the reduced Lagrangian (57)) this leads to the following current-current interaction

$$D_{\mu\nu}^R(q) J^\mu J^\nu = J_0 \frac{1}{\vec{q}^2 + M^2} J_0 + J_i J_j \left( \delta_{ij} - \frac{q_i q_j}{\vec{q}^2 + M^2} \right) \frac{1}{q^2 - M^2}. \quad (60)$$

It is the generalization of the complete radiation photon propagator (21) in QED. The propagator (60) contains the instantaneous part as a analogue of the Coulomb instantaneous one in the propagator (21).

As it was shown in [25], the Lorentz transformations of classical radiation variables coincide with the quantum ones and they both (quantum and classical) correspond to the transition to another Lorentz reference frame distinguished by another time-axis, where the relativistic covariant propagator takes the form

$$D_{\mu\nu}^R(q|n) = \frac{n_\mu n_\nu}{M_W^2 + |q^\perp|^2} - \left( \delta_{\mu\nu}^\perp - \frac{q_\mu^\perp q_\nu^\perp}{M_W^2 + |q^\perp|^2} \right) \frac{1}{q^2 - M_W^2 + i\epsilon} \quad (61)$$

with  $q_\mu^\perp = q_\mu - n_\mu(qn)$ ,  $\delta_{\mu\nu}^\perp = \delta_{\mu\nu} - n_\mu n_\nu$ . Therefore, we shall use the radiation propagator (61), where  $n_\mu$  is treated as the initial datum operator (39), instead of the conventional frame free massive vector propagator [6]

$$J^\mu D_{\mu\nu}^F(q) J^\nu = -\frac{1}{q^2 - M^2} J^\mu \left( g_{\mu\nu} - \frac{q_\mu q_\nu}{M^2} \right) J^\nu \quad (62)$$

In contrast to this conventional frame free massive vector propagator the radiation propagator (60) is regular in the limit  $M \rightarrow 0$  and is well behaved for large momenta.

For a better comparison with the conventional covariant propagator (62) we rewrite the propagator (60) in the alternative form

$$D_{\mu\nu}^R(q)J^\mu J^\nu = -\frac{1}{q^2 - M^2} \left( J_\nu^2 + \frac{(J_i q_i)^2 - (J_0 q_0)^2}{(\vec{q}^2 + M^2)} \right). \quad (63)$$

Hence we see that for massive QED, where the vector field is coupled to a conserved current ( $q_\mu J^\mu = 0$ ), we find that the effective current-current interactions mediated by the propagator of the radiation fields (60) and by the conventional frame free propagator (62) coincide

$$J^\mu D_{\mu\nu}^R J^\nu = J^\mu D_{\mu\nu}^F J^\nu. \quad (64)$$

If the current is not conserved  $J_0 q_0 \neq J_k q_k$ , the radiation field variables with the propagator (60) (or (61)) is inequivalent to the frame free variables with the propagator (62) because different variables corresponds to different *gauges of physical sources*.

In the next Section, we show the dependence of physical results on *gauge of physical sources*.

## 4 Observational tests

### 4.1 Chiral Bosonization of EW Interaction

In order to illustrate the inequivalence of Dirac's method (57) and the frame free one [6], we can consider the  $K \rightarrow \pi$  transition amplitude and the kaon decays, where the dominance of the instantaneous interactions was recently predicted [26]. Recall that the observation of kaon weak decays has been crucial for the modern theory of particle physics [27].

It was accepted to describe weak decays in the framework of electroweak (EW) theory at the quark QCD level including current vector boson weak interactions [28, 29]

$$\mathcal{L}_{(J)} = -(J_\mu^- W_\mu^+ + J_\mu^+ W_\mu^-) = -\frac{e}{2\sqrt{2}\sin\theta_W} (\underline{J}_\mu^- W_\mu^+ + \underline{J}_\mu^+ W_\mu^-), \quad (65)$$

where  $\underline{J}_\mu^+ = \bar{d}'\gamma_\mu(1 - \gamma_5)u$ ;  $\bar{d}' = d\cos\theta_C + s\sin\theta_C$ ,  $\theta_C$  is a Cabbibo angle  $\sin\theta_C = 0.223$ .

However, a consistent theory at large distances of QCD is not yet constructed up to now. Therefore, now the most effective method of analysis of kaon decay physics [30, 31, 32, 33] is the chiral perturbation theory [34, 35] (the list of arguments in favor of this perturbation theory is given in Appendix B).

The quark content of  $\pi^+$  and  $K^+$  mesons  $\pi^+ = (\bar{d}, u)$ ,  $K^+ = (\bar{s}, u)$ ,  $\bar{K}^0 = (\bar{s}, d)$  leads to the effective chiral hadron currents  $\underline{J}_\mu^\pm$  in the Lagrangian (65)

$$\underline{J}_\mu^\pm = [\underline{J}_\mu^1 \pm i\underline{J}_\mu^2] \cos\theta_C + [\underline{J}_\mu^4 \pm i\underline{J}_\mu^5] \sin\theta_C, \quad (66)$$

where using the Gell-Mann matrixes  $\lambda^k$  one can define the meson current as [34]

$$i \sum \lambda^k \underline{J}_\mu^k = i\lambda^k (V_\mu^k - A_\mu^k)^k = F_\pi^2 e^{i\xi} \partial_\mu e^{-i\xi}, \quad (67)$$

$$\xi = F_\pi^{-1} \sum_{k=1}^8 M^k \lambda^k = F_\pi^{-1} \begin{pmatrix} \pi^0 + \frac{\eta}{\sqrt{3}} & \pi^+ \sqrt{2} & K^+ \sqrt{2} \\ \pi^- \sqrt{2} & -\pi^0 + \frac{\eta}{\sqrt{3}} & K^0 \sqrt{2} \\ K^- \sqrt{2} & \bar{K}^0 \sqrt{2} & -\frac{2\eta}{\sqrt{3}} \end{pmatrix}. \quad (68)$$




Figure 1: Axial (a) and vector (b) current contribution into  $K^+ \rightarrow \pi^+$  transition

In the first orders in mesons one can write

$$V_\mu^- = \sqrt{2} [\sin \theta_C (K^- \partial_\mu \pi^0 - \pi^0 \partial_\mu K^-) + \cos \theta_C (\pi^- \partial_\mu \pi^0 - \pi^0 \partial_\mu \pi^-)] + \dots \quad (69)$$

and

$$A_\mu^- = \sqrt{2} F_\pi (\partial^\mu K^- \sin \theta_C + \partial^\mu \pi^- \cos \theta_C) + \dots; \quad (70)$$

here  $F_\pi \simeq 92$  MeV. The right form of chiral Lagrangian of the electromagnetic interaction of mesons can be constructed by the covariant derivative  $\partial_\mu \chi^\pm \rightarrow D_\mu \chi^\pm \equiv (\partial_\mu \pm ie A_\mu) \chi^\pm$ , where  $\chi^\pm = K^\pm, \pi^\pm$ .

We suppose also that the quark content of the mesons determines hadronization of QCD [20, 21] conserving its chiral and gauge symmetries.

## 4.2 $K \rightarrow \pi$ Transition Amplitude and the Rule $\Delta T = \frac{1}{2}$

Let us consider the  $K^+ \rightarrow \pi^+$  transition amplitude

$$\langle \pi^+ | -i \int dx^4 dy^4 J^\mu(x) D_{\mu\nu}^W(x-y) J^\nu(y) | K^+ \rangle = i(2\pi)^4 \delta^4(k-p) G_{EW} \Sigma(k^2), \quad (71)$$

in the first order of the EW perturbation theory in the Fermi coupling constant

$$G_{EW} = \frac{\sin \theta_C \cos \theta_C}{8M_W^2} \frac{e^2}{\sin^2 \theta_W} \equiv \sin \theta_C \cos \theta_C \frac{G_F}{\sqrt{2}}, \quad (72)$$

comparing two different W-boson field propagators, the radiation (R) propagator (61)

$$D_{\mu\nu}^W(q) \rightarrow D_{\mu\nu}^R(q|n) = \frac{n_\mu n_\nu}{M_W^2 + |q^\perp|^2} - \left( \delta_{\mu\nu}^\perp - \frac{q_\mu^\perp q_\nu^\perp}{M_W^2 + |q^\perp|^2} \right) \frac{1}{q^2 - M_W^2 + i\epsilon}, \quad (73)$$

where  $q_\mu^\perp = q_\mu - n_\mu(qn)$ ,  $\delta_{\mu\nu}^\perp = \delta_{\mu\nu} - n_\mu n_\nu$  in the frame  $n_\mu = k_\mu / \sqrt{k_\mu^2}$ , and the frame free (F) propagator (62)

$$D_{\mu\nu}^W(q) \rightarrow D_{\mu\nu}^F(q) = -\frac{1}{q^2 - M_W^2} \left( g_{\mu\nu} - \frac{q_\mu q_\nu}{M_W^2} \right). \quad (74)$$

These propagators give the expressions corresponding to the diagrams in Fig 1

$$\Sigma(k^2) \rightarrow \Sigma^R(k^2) = 2F_\pi^2 k_\mu D_{\mu\nu}^R(k|n) k_\nu + 2i \int \frac{d^4 q}{(2\pi)^4} \frac{(2k_\mu + q_\mu) D_{\mu\nu}^R(-q|n) (2k_\nu + q_\nu)}{(k+q)^2 - m_\pi^2 + i\epsilon}, \quad (75)$$

$$\Sigma(k^2) \rightarrow \Sigma^F(k^2) = 2F_\pi^2 k_\mu D_{\mu\nu}^F(k) k_\nu + 2i \int \frac{d^4 q}{(2\pi)^4} \frac{(2k_\mu + q_\mu) D_{\mu\nu}^F(-q) (2k_\nu + q_\nu)}{(k+q)^2 - m_\pi^2 + i\epsilon} = \quad (76)$$

$$= 2k_\mu^2 \frac{F_\pi^2}{M_W^2} + \int \frac{d^4 q}{(2\pi)^4} \frac{(2k_\mu + q_\mu)(2k_\nu + q_\nu)}{(M_W^2 - q^2 - i\epsilon)[(k+q)^2 - m_\pi^2 + i\epsilon]} \left[ \delta^{\mu\nu} - \frac{q^\mu q^\nu}{M_W^2} \right]. \quad (77)$$

Both versions R and F coincide in the case of the axial contribution corresponding to the first diagram in Fig. 1 and both they reduce to the instantaneous interaction contribution, because

$$k^\mu k^\nu D_{\mu\nu}^F(k) \equiv k^\mu k^\nu D_{\mu\nu}^R(k|n) = \frac{k^2}{M_W^2}.$$

However, in the case of the vector contribution corresponding to the second diagram in Fig. 1 the radiation version differs from the frame free one (77)<sup>10</sup>.

In contrast to the frame free version (77), two radiation variable diagrams shown in Fig. 1 in the frame  $n_\mu \sim k_\mu = (k_0, 0, 0, 0)$  are reduced to the instantaneous interaction contribution

$$i(2\pi)^4 \delta^4(k-p) G_{\text{EW}} \Sigma^R(k^2) = \langle \pi^+ | -i \int d^4x J_0(x) \frac{1}{\Delta - M_W^2} J_0(x) | K^+ \rangle \quad (78)$$

with the normal ordering of the pion fields, so that

$$\Sigma^R(k^2) = 2k_\mu^2 \left[ \frac{F_\pi^2}{M_W^2} + \frac{1}{(2\pi)^3} \int \frac{d^3l}{2E_\pi(\vec{l})} \frac{1}{M_W^2 + \vec{l}^2} \right]; \quad (79)$$

here  $E_\pi(\vec{l}) = \sqrt{m_\pi^2 + \vec{l}^2}$  is the energy of  $\pi$ -meson. The reduction of the radiation variable loop diagrams to the instantaneous interaction ones demonstrates that the radiation variables allow to separate the low-energy region from the high-energy one. This separation justifies the application of the low-energy chiral perturbation theory [36], where the summation of the chiral series can be considered here as the meson form factors [30, 32, 33]. Finally, the radiation variables (75) and (78) give the result

$$\Sigma^R(k^2) = 2k_\mu^2 \left[ \frac{F_\pi^2}{M_W^2} + \frac{1}{(2\pi)^3} \int \frac{d^3l}{2E_\pi(\vec{l})} \frac{f_{K\pi W}(-\vec{l}^2) f_{\pi\pi W}(-\vec{l}^2)}{M_W^2 + \vec{l}^2} \right] \equiv 2k_\mu^2 \frac{F_\pi^2}{M_W^2} g_8; \quad (80)$$

here  $f_{K\pi W}^V(-\vec{l}^2)$  and  $f_{\pi\pi W}^V(-\vec{l}^2)$  are the weak vector form factors introduced in the chiral perturbation theory [36] as the natural regularization of the low-energy meson instantaneous interactions by the next orders, and

$$g_8 = 1 + \frac{M_W^2}{(2\pi)^2 F_\pi^2} \int \frac{d|\vec{l}|}{E_\pi(\vec{l})} \frac{\vec{l}^2 f_{K\pi W}^V(-\vec{l}^2) f_{\pi\pi W}^V(-\vec{l}^2)}{M_W^2 + \vec{l}^2}, \quad (81)$$

is the parameter of the enhancement of the probability of the axial  $K^+ \rightarrow \pi^+$  transition. The relation (81), where  $g_8 = 5.1$  [31, 37], can be considered as the low-energy sum rule for meson form factors [38] in the space-like region.

This result shows that the instantaneous vector interaction can increase the axial interaction  $K^+ \rightarrow \pi^+$  transition in  $g_8$  times and give a new term describing the  $K^0 \rightarrow \pi^0$  transition proportional to  $g_8 - 1$ .

Using the covariant perturbation theory [39] developed as the series  $\underline{J}_\mu^k(\gamma \oplus \xi) = \underline{J}_\mu^k(\xi) + F_\pi^2 \partial_\mu \gamma^k + \gamma^i f_{ijk} \underline{J}_\mu^j(\xi) + O(\gamma^2)$  with respect to quantum fields  $\gamma$  added to  $\xi$  as the product

<sup>10</sup>The Faddeev equivalence theorem [3] is not valid, because the vector current  $J_\mu = K \partial_\mu \pi - \pi \partial_\mu K$  becomes the vertex  $\Gamma_\mu = K \partial_\mu D_\pi - D_\pi \partial_\mu K$ , where one of fields is replaced by its propagator  $\square D_\pi = \delta(x)$ , and  $\partial_\mu \Gamma^\mu \neq 0$ .




Figure 2: Axial current contribution

$e^{i\gamma}e^{i\xi} \equiv e^{i(\gamma \oplus \xi)}$  [39], one can see that the normal ordering  $N(\vec{x} - \vec{y}) = \langle 0 | \gamma^i(x) \gamma^{i'}(y) | 0 \rangle = \delta^{ii'} \int d^3l e^{i\vec{l} \cdot (\vec{x} - \vec{y})} / (2E_\pi(\vec{l}))$  in the product of the currents can lead to an effective Lagrangian

$$\int d^3z \frac{N(\vec{z}) e^{-M_W |\vec{z}|}}{4\pi |\vec{z}|} \underline{J}_\mu^j(\xi(x)) \underline{J}_\mu^{j'}(\xi(z+x)) [(f_{ij1} + i f_{ij2})(f_{ij'4} - i f_{ij'5}) + h.c.] \langle 0 | \gamma^i(x) \gamma^{i'}(y) | 0 \rangle.$$

In the limit  $M_W \rightarrow \infty$ , one can neglect the dependence of the current on  $\vec{z}$ , and we get the effective Lagrangians [40]

$$\mathcal{L}_{(\Delta T = \frac{1}{2})} = \frac{G_F}{\sqrt{2}} g_8 \cos \theta_C \sin \theta_C \left[ (\underline{J}_\mu^1 + i \underline{J}_\mu^2)(\underline{J}_\mu^4 - i \underline{J}_\mu^5) - (\underline{J}_\mu^3 + \frac{1}{\sqrt{3}} \underline{J}_\mu^8)(\underline{J}_\mu^6 - i \underline{J}_\mu^7) + h.c. \right], \quad (82)$$

$$\mathcal{L}_{(\Delta T = \frac{3}{2})} = \frac{G_F}{\sqrt{2}} \cos \theta_C \sin \theta_C \left[ (\underline{J}_\mu^3 + \frac{1}{\sqrt{3}} \underline{J}_\mu^8)(\underline{J}_\mu^6 - i \underline{J}_\mu^7) + h.c. \right]. \quad (83)$$

This Lagrangian at the level of the tree diagrams describes the nonleptonic decays in satisfactory agreement with experimental data within the accuracy  $20 \div 30\%$  [34, 40].

### 4.3 The $K^+ \rightarrow \pi^+ + l + \bar{l}$ amplitude

The result of calculation of the axial contributions to the amplitude of the process  $K^+ \rightarrow \pi^+ + l + \bar{l}$  within the framework of chiral Lagrangian (65), (66) including phenomenological meson form factors shown in Fig. 2 as a fat dot takes the form

$$T_{(K^+ \rightarrow \pi^+ l^- \bar{l}^-)}^{AA} = 2e G_{EW} L_\nu D_{\mu\nu}^{\gamma(rad)}(q)(k_\mu + p_\mu) t^{AA}(q^2, k^2, p^2), \quad (84)$$

where  $G_{EW}$  is the constant (72),  $L_\mu = \bar{l} \gamma_\mu l$  is leptonic current,

$$t^{AA}(q^2, k^2, p^2) = F_\pi^2 \left[ \frac{f_\pi^V(q^2) k^2}{m_\pi^2 - k^2 - i\epsilon} + \frac{f_K^V(q^2) p^2}{M_K^2 - p^2 - i\epsilon} + \frac{f_K^A(q^2) + f_\pi^A(q^2)}{2} \right], \quad (85)$$

and  $f_{\pi,K}^{(A,V)}(q^2)$  are meson form factors. In this expression the propagator  $D_{\mu\nu}^R(p|n)$  given by Eq. (61) keeps only the covariant instantaneous part  $\frac{p_\mu p_\nu}{p^2 M_W^2}$ .




Figure 3: Vector current contribution

On the mass-shell the sum (85) takes the form

$$\begin{aligned}
 t^{AA}(q^2, M_K^2, m_\pi^2) &= \\
 = t(q^2) &= F_\pi^2 \left[ \frac{f_K^A(q^2) + f_\pi^A(q^2)}{2} - f_\pi^V(q^2) + [f_K^V(q^2) - f_\pi^V(q^2)] \frac{m_\pi^2}{M_K^2 - m_\pi^2} \right].
 \end{aligned} \tag{86}$$

As can be seen in [30, 31], the amplitude for  $K^+ \rightarrow \pi^+ + l + \bar{l}$  vanishes at the tree level, where form factors are equal to unit  $t(q^2)|_{f^V=f^A=1} = 0$ . It is well known that these form factors in the limit  $q^2 \rightarrow 0$  take the form  $f_{\pi,K}^{A,V}(q^2) = 1 - \frac{q^2}{6} < r^2 >_{\pi,K}^{(A,V)}$ , where  $< r^2 >_{\pi,K}^{(A,V)}$  are the axial and vector mean square radii of mesons, respectively, in agreement with the chiral perturbation theory [30, 34, 36].

The result of calculation of vector contributions to the amplitude of the process  $K^+ \rightarrow \pi^+ + l + \bar{l}$  within the framework of chiral Lagrangian (65), (12) including phenomenological meson form factors shown in Fig. 3 as a fat dot takes the form

$$T_{(K^+ \rightarrow \pi^+ l \bar{l})}^{VV(\text{rad})} = 2eG_{\text{EW}} L_\nu D_{\mu\nu}^{\gamma(\text{rad})}(q)(k_\mu + p_\mu) t^{VV}(q^2, k^2, p^2), \tag{87}$$

where we use the propagator (61) in the initial datum frame (39)

$$\begin{aligned}
 t^{VV}(q^2, k^2, p^2) &= \frac{1}{2(2\pi)^2} \int \frac{d\vec{l}}{E_\pi(\vec{l})} \frac{\vec{l}^2}{E_\pi(\vec{l})} \left\{ f_{K\pi W}^V(\vec{l}^2) f_{\pi\pi W}^V(\vec{l}^2) \left[ \frac{f_\pi^V(q^2) k^2}{m_\pi^2 - k^2 - i\epsilon} + \right. \right. \\
 &\quad \left. \left. + \frac{f_K^V(q^2) p^2}{M_K^2 - p^2 - i\epsilon} \right] + \frac{f_\pi^A(q^2) f_{K\pi W}^V(\vec{l}^2) f_{\pi\pi W\gamma}^V(\vec{l}^2) + f_K^A(q^2) f_{\pi\pi W}^V(\vec{l}^2) f_{K\pi W\gamma}^V(\vec{l}^2)}{2} \right\};
 \end{aligned}$$

here  $f_{\pi\pi W\gamma}^V(\vec{l}^2)$ ,  $f_{K\pi W\gamma}^V(\vec{l}^2)$  are the four particle interaction form factors which should coincide with the three particle interaction form factors  $f_{\pi\pi W\gamma}^V = f_{\pi\pi W}^V$ ,  $f_{K\pi W\gamma}^V = f_{K\pi W}^V$  in accordance with the gauge invariance of the strong interactions.

Using the result (81) one can write on the mass shell  $k^2 = M_K^2, p^2 = m_\pi^2$ :

$$t^{VV}(q^2, M_K^2, m_\pi^2) = (g_8 - 1)t(q^2), \tag{88}$$

$$t^{AA}(q^2, M_K^2, m_\pi^2) + t^{VV}(q^2, M_K^2, m_\pi^2) = g_8 t(q^2), \tag{89}$$



where  $g_8, t(q^2)$  are given by Eqs. (81) and (86), respectively.

Finally, the result of calculation of axial and vector contributions to the amplitude of the process  $K^+ \rightarrow \pi^+ + l + \bar{l}$  within the framework of chiral Lagrangian (66),(69) including phenomenological meson form factors takes the form

$$T_{(K^+ \rightarrow \pi^+ l + \bar{l})}^{(\text{rad})} = 2eg_8 G_{\text{EW}} L_\nu D_{\mu\nu}^{\gamma(\text{rad})}(q)(k_\mu + p_\mu) t(q^2). \quad (90)$$

This amplitude leads to the total decay rate for the transition  $K^+ \rightarrow \pi^+ e^+ e^-$

$$\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)}((M_K - m_\pi)^2) = \bar{\Gamma}_{e^+ e^-} \int_{4m_e^2}^{(M_K - m_\pi)^2} \frac{dq^2}{M_K^2} \rho(q^2) F(q^2), \quad (91)$$

where

$$\bar{\Gamma}_{e^+ e^-} = \frac{(s_1 c_1 c_3)^2 g_8^2 G_F^2 \alpha^2 M_K^5}{(4\pi)^4 24\pi} \Big|_{g_8=5.1} = 1.37 \times 10^{-22} \text{GeV}, \quad (92)$$

$$\begin{aligned} F(q^2) &= \left[ \frac{(4\pi)^2 t(q^2)}{q^2} \right]^2 = \\ &= \left[ \frac{(4\pi F_\pi)^2}{q^2} \right]^2 \left[ \frac{f_K^A(q^2) + f_\pi^A(q^2)}{2} - f_\pi^V(q^2) + [f_K^V(q^2) - f_\pi^V(q^2)] \frac{m_\pi^2}{M_K^2 - m_\pi^2} \right]^2 \end{aligned} \quad (93)$$

and

$$\rho(q^2) = \left( 1 - \frac{4m_l^2}{q^2} \right)^{1/2} \times \left( 1 + \frac{2m_l^2}{q^2} \right) \lambda^{3/2}(1, q^2/M_K^2, m_\pi^2/M_K^2);$$

here  $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2(ab + bc + ca)$ ,  $s_1 c_1 c_3$  is the product of Cabibbo-Kobayashi-Maskawa matrix elements  $V_{ud} V_{us}$ .

The mechanism of enhancement  $|\Delta T| = \frac{1}{2}$  considered in this paper can be generalized to a description of other processes including  $K^+ \rightarrow \pi^+ \nu \bar{\nu}$  by replacing the  $\gamma$ -propagator by the Z-boson one, so that we have the relations

$$\begin{aligned} &\left[ \frac{q^2}{(4\pi F_\pi)^2} \right]^2 \frac{M_K^2}{\rho(q^2) \bar{\Gamma}_{e^+ e^-}} \frac{d\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)}(q^2)}{dq^2} = \\ &= \left[ \frac{M_Z^2}{(4\pi F_\pi)^2} \right]^2 \frac{M_K^2}{\rho(q^2) \bar{\Gamma}_{\nu \bar{\nu}}} \frac{d\Gamma_{(K^+ \rightarrow \pi^+ \nu \bar{\nu})}(q^2)}{dq^2} = \\ &= \left[ \frac{f_K^A(q^2) + f_\pi^A(q^2)}{2} - f_\pi^V(q^2) + [f_K^V(q^2) - f_\pi^V(q^2)] \frac{m_\pi^2}{M_K^2 - m_\pi^2} \right]^2. \end{aligned} \quad (94)$$

Thus, exploiting the instantaneous weak interaction mechanism of enhancement in the  $K \rightarrow \pi$  transition probability and QCD symmetry we derive the sum rule of EW vector meson form factors given by Eqs. (81), (86), (91) and their relation to the differential  $K \rightarrow \pi e^+ e^-$  decay rate (94).

#### 4.4 The form factor probe of the differential $K \rightarrow \pi e^+ e^-$ decay rate

The estimation of the meson loop contribution was made in [31] where a function  $\hat{\phi}^2(q^2)$  was used instead of the form factor rate  $F(q^2)$  (93)

$$F(q^2) \implies \hat{\phi}^2(q^2)$$

It was shown that the values  $g_8 = 5.1$ ,  $\hat{\phi}(0) = 1$  gave the total decay rate  $\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)} = 1.91 \times 10^{-23}$  GeV in the satisfactory agreement with the experimental data  $\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)} = 1.44 \pm 0.27 \times 10^{-23}$  GeV [37, 41]. However, the main contribution goes from the baryon loops [30, 32, 33]. Therefore, we discuss here the value of the differential  $K \rightarrow \pi e^+ e^-$  decay rate (93) in the chiral perturbation theory [34, 36] where both the pion loop contribution  $\Pi_\pi$  and baryon ones lead to the meson form factors and resonances [30, 36] in the Padé-type approximation

$$f_\pi^V(q^2) = 1 + M_\rho^{-2} q^2 + \alpha_0 \Pi_\pi(q^2) + \dots \simeq \frac{1}{1 - M_\rho^{-2} q^2 - \alpha_0 \Pi_\pi(q^2)} \quad (95)$$

$$f_\pi^A(q^2) \simeq f_K^A(q^2) = 1 + M_a^{-2} q^2 + \dots \simeq \frac{1}{1 - M_a^{-2} q^2} \quad (96)$$

where

$$\alpha_0 = \frac{m_\pi^2}{(4\pi F_\pi)^2} \frac{4}{3} \simeq \frac{2}{103}, \quad (97)$$

$$\Pi_\pi(t) = (1 - \bar{t}) \left( \frac{1}{\bar{t}} - 1 \right)^{1/2} \arctan \left( \frac{\bar{t}^{1/2}}{(1 - \bar{t})^{1/2}} \right) - 1; \quad 0 < t < (2m_\pi)^2 \quad (98)$$

$$\Pi_\pi(t) = \frac{\bar{t} - 1}{2} \left( 1 - \frac{1}{\bar{t}} \right)^{1/2} \left\{ i\pi - \log \frac{\bar{t}^{1/2} + (\bar{t} - 1)^{1/2}}{\bar{t}^{1/2} - (\bar{t} - 1)^{1/2}} \right\} + 1; \quad (2m_\pi)^2 < t \quad (99)$$

is the pion loop contribution [34, 36], and  $M_\rho = 771$  MeV and  $M_a = 980$  MeV are the values of resonance masses [37] that can be determined by the baryon loops [30, 32, 33, 36].

The constant approximation of the type of [31]

$$F(q^2 = 0) = F(0) = (4\pi F_\pi)^4 \frac{[f_\pi^A(q^2) - f_\pi^V(q^2)]^2}{(q^2)^2} \Big|_{q^2=0} = (4\pi F_\pi)^4 [M_\rho^{-2} - M_a^{-2}]^2 = 0.74 \quad (100)$$

corresponds to the value of  $\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)} = 1.41 \times 10^{-23}$  GeV in satisfactory agreement with the experimental data  $\Gamma_{(K^+ \rightarrow \pi^+ e^+ e^-)} = 1.44 \pm 0.27 \times 10^{-23}$  GeV [37, 41].

The form factors (95) and (96) below the two particle threshold  $q^2 < 4m_\pi^2$  determines the differential rate  $F(q^2)$  as a function of  $q^2$

$$F(q^2) = \frac{F(0)}{[1 - M_\rho^{-2} q^2]^2 [1 - M_a^{-2} q^2]^2} \longrightarrow \frac{d}{dq^2} \log F(q^2) \Big|_{q^2=0} \simeq 2[M_\rho^{-2} + M_a^{-2}] \simeq 5 \text{GeV}^{-2} \quad (101)$$

At the region above the two particle threshold  $4m_\pi^2 < q^2 < (M_K - m_\pi)^2$  there is a jump of the differential rate  $F(q^2)$  at the level of  $5 \div 10$  %

$$F(q^2) = (4\pi F_\pi)^4 \frac{[M_\rho^{-2} - M_a^{-2}]^2 + \alpha_0^2 \pi^2 (q^2 - 4m_\pi^2)/(4q^6)}{[1 - M_\rho^{-2} q^2]^2 [1 - M_a^{-2} q^2]^2} \quad (102)$$

These results (100), (101), and (102) are arguments that the detailed investigation of the differential  $K \rightarrow \pi e^+ e^-$  decay rate in the NA48/2 CERN experiment can give us information about the meson form factors. To make much more realistic analysis, we have to use the unitary and analytic model for meson form factors [38] describing very well all experimental data.

## 5 Conclusion

The instantaneous interactions are inevitable consequence of the Dirac approach to gauge theories. Recall that Dirac eliminated all fields with zero momenta (and possible negative contributions into the energy of the system) by resolution of the Gauss constraint and its diagonalization, in order to have a physical vacuum as a state with minimal energy in a comoving *inertial frame*.

This elimination unambiguously leads to the *radiation variables* associated in QED with the Coulomb *gauge constraint*. However, in SM, the *radiation variables* of the massive vector bosons do not correspond to any whatsoever *gauge constraint*. The transition from the *radiation variables* with the instantaneous interactions to a frame free formulation [6] without the instantaneous interactions is fulfilled by *gauge of physical sources*. The change of the *gauge of physical sources* is just the point where the accepted frame free methods [6] lose all instantaneous interactions together with their physical effects, including the instantaneous bound states. The strong dependence of physical results on the *gauge of physical sources* does not mean the violation of the gauge-invariance principle. This means that *frame free* method has a region of validity restricted by the *initial datum free* processes of the type of the elementary particles scattering ones [3], as it follows from the physical meaning of the concept of the *frame* revealed by its full name *frame of reference to initial data*.

In order to keep relativistic invariance of S-matrix elements between the instantaneous bound states, the instantaneous interactions are determined in an arbitrary frame, where its time-axis is treated as the operator of *initial data* acting in the complete set of the all asymptotic physical states including the bound states considered as irreducible representations of the Poincaré group [9].

In the paper, we compare this Dirac operator approach to SM with the acceptable frame free method [6] using as example the  $K^+ \rightarrow \pi^+$  transition and the semileptonic and nonleptonic kaon decay probability amplitudes. This comparison shows us that the Dirac approach to SM separates the low-energy contributions from the high-energy ones and expresses these amplitudes in terms of the hadron electroweak form factors. Therefore, the obtained amplitudes allow to extract information about form factors of  $\pi$  and  $K$  mesons from the  $K^+ \rightarrow \pi^+$  transition in the  $K^+ \rightarrow \pi^+ e^+ e^- (\mu^+ \mu^-, \nu \bar{\nu})$  and  $K^+ \pi^- \rightarrow e^+ e^- (\mu^+ \mu^-, \nu \bar{\nu})$  processes. The parameters of the differential  $K \rightarrow \pi e^+ e^-$  decay rate in the NA48/2 CERN experiment were estimated in the chiral perturbation theory.

These results show that the Dirac formulation of Standard Model can reveal new physical effects in comparison with the frame free formulation [6] used now for describing observational data. Therefore, the problem of the Dirac formulation of SM becomes topical in the light of future precision experiments.

## Acknowledgments

The authors are grateful to B.M. Barbashov, D.Yu. Bardin, A.Di Giacomo, S.B. Gerasimov, G.V. Efimov, A.V. Efremov, V.D. Kekelidze, E.A. Kuraev, V.B. Priezzhev, A.N. Sissakian, S.I. Vinitzky and M.K. Volkov for fruitful discussions. The work was in part supported by the Slovak Grant Agency for Sciences VEGA, Gr.No.2/4099/26.

# Appendix A: S-matrix for bound states in electrodynamics

## A.1. Spectral problem

Let us keep in the QED action only the instantaneous interaction (18) neglecting radiation corrections

$$S_{\text{QED}}^{(\text{rad})}[\psi, \bar{\psi}] = \int d^4x [\bar{\psi}(x)(i\partial - m)\psi(x) + \frac{1}{2} \int d^4y (\psi(y)\bar{\psi}(x))\mathcal{K}(x, y | n)(\psi(x)\bar{\psi}(y))]; \quad (\text{A.1})$$

here

$$\mathcal{K}(x, y | n) = \not{n}^{\text{cf}} V(z^\perp) \delta(z \cdot n^{\text{cf}}) \not{n}^{\text{cf}}, (\not{n} = n^\mu \gamma_\mu = \gamma \cdot n) \quad (\text{A.2})$$

is the instantaneous kernel depending on the relative coordinates  $z = x - y$  and  $V(z^\perp) = -\frac{e^2}{4\pi|z^\perp|}$  depends on the transverse components  $z_\mu^\perp = z_\mu - n_\mu^{\text{cf}}(z \cdot n^{\text{cf}})$  of the relative coordinate with respect to the time axis  $n_\mu^{\text{cf}} = (1, 0, 0, 0)$ .

It seems that a most straightforward way for constructing S-matrix of bound states in gauge theories is the redefinition of action (A.1) in terms of bilocal fields by means of the Legendre transformation [42] of the current – current interaction with a kernel  $\mathcal{K}(x, y|n)$

$$\begin{aligned} & \frac{1}{2} \int d^4x d^4y (\psi(y)\bar{\psi}(x))\mathcal{K}(x, y|n)(\psi(x)\bar{\psi}(y)) = \\ & = - \frac{1}{2} \int d^4x d^4y \mathcal{M}(x, y)\mathcal{K}^{-1}(x, y|n)\mathcal{M}(x, y) + \int d^4x d^4y (\psi(x)\bar{\psi}(y))\mathcal{M}(x, y). \end{aligned} \quad (\text{A.3})$$

and the Gauss functional integral over the bilocal fields in the generating functional

$$Z[\eta^{(\text{rad})}, \bar{\eta}^{(\text{rad})}] = \int D\mathcal{M} D\psi D\bar{\psi} \exp \left\{ (\psi\bar{\psi}, (-G_0^{-1} + \mathcal{M})) - \frac{1}{2}(\mathcal{M}, \mathcal{K}^{-1}\mathcal{M}) + i(\bar{\psi}\eta^{(\text{rad})}) + (\bar{\eta}^{(\text{rad})}\psi) \right\}; \quad (\text{A.4})$$

here we used the short - hand notation

$$\int d^4x d^4y \psi(y)\bar{\psi}(x)(i\partial - m^0)\delta^{(4)}(x - y) = (\psi\bar{\psi}, -G_0^{-1}), \quad (\text{A.5})$$

$$\int d^4x d^4y (\psi(x)\bar{\psi}(y))\mathcal{M}(x, y) = (\psi\bar{\psi}, \mathcal{M}). \quad (\text{A.6})$$

The integration over the fermi fields  $\psi, \bar{\psi}$  in (A.4) leads to the generating functional

$$Z_{FP}[\eta^{(\text{rad})}, \bar{\eta}^{(\text{rad})}] = \int D\mathcal{M} \exp \left\{ iW_{eff}[\mathcal{M}] + (\bar{\eta}^{(\text{rad})}\eta^{(\text{rad})}, (-G_0^{-1} + \mathcal{M})^{-1}) \right\} \quad (\text{A.7})$$

with the effective action [42]

$$W_{eff}[\mathcal{M}] = -\frac{1}{2}(\mathcal{M}, \mathcal{K}^{-1}\mathcal{M}) + iTr \log(-G_0^{-1} + \mathcal{M}). \quad (\text{A.8})$$

The first step to the quantization of the effective action is the determination of its minimum

$$\frac{\delta W_{eff}(\mathcal{M})}{\delta \mathcal{M}} \equiv -\mathcal{K}^{-1}\mathcal{M} + \frac{i}{G_0^{-1} - \mathcal{M}}|_{\mathcal{M}=\Sigma} = 0 \quad (\text{A.9})$$

known as the Schwinger – Dyson equation of the fermion Green function

$$\underline{G}_\Sigma(q) = \frac{1}{\not{q} - m_0 - \underline{\Sigma}(q^\perp)}, \quad (\text{A.10})$$

we denote the corresponding classical solution for the bilocal field by  $\Sigma(z) = \int d^4q \Sigma(q^\perp) e^{iq \cdot z} \delta(z \cdot n)$ .

The next step is the expansion of the effective action around the point of minimum  $\mathcal{M} = \Sigma + \widetilde{\mathcal{M}}$ ,

$$W_{eff}(\Sigma + \widetilde{\mathcal{M}}) = W_{eff}^{(2)} + W_{int}; \quad (\text{A.11})$$

$$W_{eff}^{(2)}(\widetilde{\mathcal{M}}) = W_Q(\Sigma) + \left[ -\frac{1}{2}(\widetilde{\mathcal{M}}, \mathcal{K}^{-1} \widetilde{\mathcal{M}}) + \frac{i}{2} (G_\Sigma \widetilde{\mathcal{M}})^2 \right]; \quad (\text{A.12})$$

$$W_{int} = \sum_{n=3}^{\infty} W^{(n)} = i \sum_{n=3}^{\infty} \frac{1}{n} (G_\Sigma \widetilde{\mathcal{M}})^n, \quad (G_\Sigma = (G_0^{-1} - \Sigma)^{-1}), \quad (\text{A.13})$$

where the small fluctuations of the bilocal field  $\widetilde{\mathcal{M}}(x, y) = \widetilde{\mathcal{M}}(z = x - y | X = (x + y)/2)$  in the effective action (A.11) can be presented as the series of a complete set of the solutions  $\Gamma$  of the classical equation (known as the Bethe – Salpeter one)

$$\frac{\delta^2 W_{eff}(\Sigma + \widetilde{\mathcal{M}})}{\delta \widetilde{\mathcal{M}}^2} \Big|_{\widetilde{\mathcal{M}}=0} \cdot \Gamma = 0 \quad (\text{A.14})$$

with a set of quantum numbers ( $A$ ) including masses  $M_A = \sqrt{\mathcal{P}_\mu^2}$  and energies  $\omega_A = \sqrt{\vec{\mathcal{P}}^2 + M_A^2}$ . In the momentum representation this series takes the form

$$\widetilde{\mathcal{M}}(z | X) = \sum_A \int \frac{d^3 \vec{\mathcal{P}}}{(2\pi)^{3/2} \sqrt{2\omega_A}} \frac{d^4 q e^{iz \cdot q}}{(2\pi)^4} \{ e^{i\mathcal{P} \cdot X} \Gamma_A(q^\perp | \mathcal{P}) a_A^+(\mathcal{P}) + e^{-i\mathcal{P} \cdot X} \bar{\Gamma}_A(q^\perp | -\mathcal{P}) a_A^-(\mathcal{P}) \}, \quad (\text{A.15})$$

where  $\widetilde{\mathcal{M}}(z | X) \sim \delta(z \cdot n)$  and  $a_A^+, a_{A'}^-$  are the bound state creation and annihilation operators that obey the commutation relations

$$\left[ a_{A'}^-(\vec{\mathcal{P}}'), a_A^+(\vec{\mathcal{P}}) \right] = \delta_{A'A} \delta^3(\vec{\mathcal{P}}' - \mathcal{P}), \quad \left[ a_A^\pm(\mathcal{P}), a_{A'}^\pm(\mathcal{P}') \right] = 0. \quad (\text{A.16})$$

In the momentum representation  $\underline{\Gamma}(k^\perp, \mathcal{P})$ , Eq. (A.14) takes the form

$$\underline{\Gamma}(k^\perp, \mathcal{P}) = i \int \frac{d^4 q}{(2\pi)^4} \underline{V}(k^\perp - q^\perp) \not{n} \left[ \underline{G}_{\Sigma a}(q + \mathcal{P}/2) \Gamma(q^\perp, \mathcal{P}) \underline{G}_{\Sigma b}(q - \mathcal{P}/2) \right] \not{n}, \quad (\text{A.17})$$

where  $G_{\Sigma a}$ ,  $G_{\Sigma b}$  are the fermion propagators of the two particles ( $a$ ) and ( $b$ ). Solutions of equation (A.14) satisfy the normalization condition [14]

$$i \frac{d}{d\mathcal{P}_0} \int \frac{d^4 q}{(2\pi)^4} \text{tr} \left[ \underline{G}_\Sigma(q - \frac{\mathcal{P}}{2}) \bar{\Gamma}_A(q^\perp | -\mathcal{P}) \underline{G}_\Sigma(q + \frac{\mathcal{P}}{2}) \Gamma_A(q^\perp | \mathcal{P}) \right] = 2\omega_A \quad (\text{A.18})$$

The corresponding Green function of the bilocal field takes form

$$\mathcal{G}(q^\perp, p^\perp | \mathcal{P}) = \sum_H \left\{ \frac{\Gamma_H(q^\perp | \mathcal{P}) \bar{\Gamma}_H(p^\perp | -\mathcal{P})}{(\mathcal{P}_0 - \omega_H - i\varepsilon) 2\omega_H} - \frac{\Gamma_H(q^\perp | -\mathcal{P}) \bar{\Gamma}_H(p^\perp | \mathcal{P})}{(\mathcal{P}_0 - \omega_H - i\varepsilon) 2\omega_H} \right\}. \quad (\text{A.19})$$

The Bethe – Salpeter field  $\Gamma(q^\perp, \mathcal{P})$  is connected with the wave function by the relation (see [20, 21])

$$\Phi_{ab}(q^\perp, \mathcal{P}) = \frac{i}{2\pi} \int \left[ \frac{\mathcal{P} \cdot dq}{M_A} \right] \left[ \underline{G}_{\Sigma a}(q + \mathcal{P}/2) \Gamma(q^\perp, \mathcal{P}) \underline{G}_{\Sigma b}(q - \mathcal{P}/2) \right] \quad (\text{A.20})$$

In the nonrelativistic limit in the frame  $n = n^{(cf)} = (1, 0, 0, 0)$  this wave function reduces to the Schrödinger one (A.22)

$$\Phi_A^{\alpha\beta}(\vec{q}, M_A) = \left( \frac{1 + \gamma_0}{2} \gamma_5 \right)^{\alpha\beta} \underline{\Psi}_{Sch}(\vec{q}) \sqrt{\frac{M_A}{4}} \quad (\text{A.21})$$

satisfying the standard Schrödinger equation

$$\left(-\frac{1}{2\mu}\frac{d^2}{dz^2} + V(\vec{z})\right)\underline{\Psi}_{Sch}(\vec{z}) = \epsilon\underline{\Psi}_{Sch}(\vec{z}) \quad (\text{A.22})$$

where  $\underline{\Psi}_{Sch}(\vec{z}) = \int \frac{d^3q}{(2\pi)^3} e^{i\vec{q}\vec{z}} \Psi_{Sch}(\vec{q})$  is a normalizable wave function ( $\int d^3z \|\underline{\Psi}_{Sch}(z)\|^2 = 1$ ) in a comoving frame  $\mathcal{P}_\mu = (M_A, 0, 0, 0)$ ; here  $M_A = (M_p + m_e - \epsilon)$  is the mass of an atom,  $M_p, m_e$  are masses of proton and electron,  $\mu = M_p \cdot m_e / (M_p + m_e)$  is the reduced mass, and  $z_k = (x - y)_k$  are relative coordinates.

## A.2. S-matrix elements

The next step is the construction of the relativistic covariant S-matrix elements considered as probability amplitude transitions between the all asymptotic states. These states includes bound states given by the spectral equation (36) and all Lorentz transformations of the comoving frame distinguished by the time-axis  $n^{(\text{cf})}$ . The Lorentz transformations of the time-axis  $n^{(\text{cf})} \rightarrow n^{(\text{ID})}$  mean the changes of the initial data (ID) of the bound states, i.e. their total momenta  $\mathcal{P}_{A\mu} = M_A n_\mu^{(\text{ID})}$ . These bound states correspond to the instantaneous interaction with a new time-axis  $n^{(\text{ID})}$

$$\mathcal{K}(x, y | n^{(\text{ID})}) = \not{n}^{(\text{ID})} V(z_\mu - n_\mu^{(\text{ID})} z \cdot n^{(\text{ID})}) \delta(z \cdot n^{(\text{ID})}) \not{n}^{(\text{ID})}. \quad (\text{A.23})$$

One can see that in the space of the bilocal fields (A.15) this unit time-axis  $n^{(\text{ID})}$  is proportional to an eigenvalue of the total momentum “operator” (i.e. the derivative with respect to total coordinate)

$$n_\mu^{(\text{ID})} \mathcal{M}(z|X) \sim \frac{1}{i} \frac{\partial}{\partial X_\mu} \mathcal{M}(z|X) \quad (\text{A.24})$$

in agreement with the Markov – Yukawa definition of the relativistic simultaneity in terms of bilocal fields  $\mathcal{M}(z|X) \equiv \mathcal{M}(x, y)$  [43]

$$z_\mu \frac{\partial}{\partial X_\mu} \mathcal{M}(z|X) = 0. \quad (\text{A.25})$$

Recall that this Markov – Yukawa definition [43] of the relativistic simultaneity in terms of bilocal fields has a deeper mathematical meaning as a constraint of irreducible nonlocal representations of the Poincaré group [9] for an arbitrary bilocal field  $\mathcal{M}(x, y) = \mathcal{M}(z|X)$  [20, 44].

The relativistic covariant constraint-shell quantization gives both the spectrum of bound states and their S-matrix elements.

Using the decomposition over the bound state quantum numbers ( $H$ )

$$\tilde{\Phi}_{H(ab)}(q^\perp | \mathcal{P}) = G_{\Sigma a}(q + \mathcal{P}/2) \Gamma_{H(ab)}(q^\perp | \mathcal{P}), \quad (\text{A.26})$$

we can write the S-matrix elements for the interaction  $W^{(n)}$  (A.13) between the vacuum and  $n$  bound states

$$\langle H_1 \mathcal{P}_1, \dots, H_n \mathcal{P}_n | iW^{(n)} | 0 \rangle = -i(2\pi)^4 \delta^4 \left( \sum_{i=1}^n \mathcal{P}_i \right) \prod_{j=1}^n \left[ \frac{1}{(2\pi)^3 2\omega_j} \right]^{1/2} M^{(n)}(\mathcal{P}_1, \dots, \mathcal{P}_n) \quad (\text{A.27})$$

$$\begin{aligned} M^{(n)} = & \int \frac{id^4 q}{(2\pi)^4 n} \sum_{\{i_k\}} \tilde{\Phi}_{H_{i_1}}^{a_1, a_2}(q | \mathcal{P}_{i_1}) \tilde{\Phi}_{H_{i_2}}^{a_2, a_3}(q - \frac{\mathcal{P}_{i_1} + \mathcal{P}_{i_2}}{2} | \mathcal{P}_{i_2}) \tilde{\Phi}_{H_{i_3}}^{a_3, a_4} \left( q - \frac{2\mathcal{P}_{i_2} + \mathcal{P}_{i_1} + \mathcal{P}_{i_3}}{2} | \mathcal{P}_{i_3} \right) \\ & \dots \tilde{\Phi}_{H_{i_n}}^{a_n, a_1} \left( q - \frac{2(\mathcal{P}_{i_2} + \dots + \mathcal{P}_{i_{n-1}}) + \mathcal{P}_{i_1} + \mathcal{P}_{i_n}}{2} | \mathcal{P}_{i_n} \right), \end{aligned}$$

where  $\{i_k\}$  denotes permutations over  $i_k$ ; the amplitude  $\langle H_1 \mathcal{P}_1, \dots, H_l \mathcal{P}_l | i W^{(n)} | H_{l+1} \mathcal{P}_{l+1}, \dots, H_n \mathcal{P}_n \rangle$  can be obtained by the change of the momentum signs  $\mathcal{P}_{l+1}, \dots, \mathcal{P}_n \rightarrow -\mathcal{P}_{l+1}, \dots, -\mathcal{P}_n$ .

Expressions (A.15), (A.19), (A.26), (A.27) represents Feynman rules for the construction of a quantum field theory with the action (A.11) in terms of bilocal fields.

It was shown [45] that this quantum field theory with a separable potential of the instantaneous interaction leads to the well-known Nambu Jona-Lasinio model [48, 49] and the phenomenological chiral Lagrangians [36] used for the description of the low-energy meson physics.

In the context of the Dirac approach to gauge theory, to solve the problem of hadronization in QCD, one needs to answer the following questions: i) What is the origin of the “separable potential” of hadronization in the non-Abelian theory? ii) How to combine the Schrödinger equation for heavy quarkonia (that is derived by the residuum of poles of the quark Green functions) with the quark confinement? iii) What is the origin of the additional mass of the ninth pseudoscalar meson?

## Appendix B: Instantaneous interactions in QCD

### B.1. QCD action and monopole vacuum

QCD was proposed [50] as the non-Abelian  $SU_c(3)$  theory with the action functional

$$W = \int d^4x \left\{ -\frac{1}{4} F_{\mu\nu}^a{}^2 + \bar{\psi} [i\gamma^\mu (\partial_\mu + \hat{A}_\mu) - m] \psi \right\}, \quad (\text{B.1})$$

where  $\psi$  and  $\bar{\psi}$  are the fermion quark fields,  $\hat{A}_\mu = g \frac{\lambda_a}{2i} A_\mu^a$ ,

$$F_{0i}^a = \partial_0 A_i^a - D_i^{ab}(A) A_0^b, \quad F_{jk}^a = \partial_j A_k^a - \partial_k A_j^a + g f^{abc} A_j^b A_k^c = \epsilon_{ijk} B_i^a \quad (\text{B.2})$$

are non-Abelian gluon electric and magnetic tensions, and

$$D_i^{ab}(A) = \delta^{ab} \partial_i + g f^{acb} A_i^c \quad (\text{B.3})$$

is the covariant derivative. The action (B.1) is invariant with respect to gauge transformations  $u(t, \vec{x})$

$$\hat{A}_\mu^u(x_0, \vec{x}) = u(x_0, \vec{x}) \left( \hat{A}_\mu + \partial_\mu \right) u^{-1}(x_0, \vec{x}), \quad \psi^u = u(x_0, \vec{x}) \psi. \quad (\text{B.4})$$

The class of functions with finite energy density includes the monopole - type functions  $A_i \sim O(1/r)$  like the Coulomb potential.

The asymptotic freedom phenomenon is considered as one of the arguments in favor of that the standard perturbation theory in QCD is unstable [55, 56]. One of candidates of the stable vacua in the Minkowskian space-time is the stationary monopole - type solution of the classical equation

$$\hat{A}_0^{\text{vacuum}}(x_0, \vec{x}) = 0, \quad \hat{A}_k^{\text{vacuum}}(x_0, \vec{x}) = \hat{\Phi}_k(0, \vec{x}) = -i \frac{\lambda_A^a}{2} \epsilon_{iak} \frac{x^k}{r^2} f(r), \quad (\text{B.5})$$

where the antisymmetric  $SU(3)$  matrices  $\lambda_A^a$  are denoted by  $\lambda_A^1 := \lambda^2$ ,  $\lambda_A^2 := \lambda^5$ ,  $\lambda_A^3 := \lambda^7$ , and  $r = |\vec{x}|$ , the vacuum field contains only one function  $f(r)$ .

In the following we discuss the  $SU(2)$  case  $\lambda_A^a \rightarrow \tau^a$ , ( $a = 1, 2, 3$ ) [21]. In this case, the classical equation for the function  $f$  in Eq. (B.5) takes the form

$$D_k^{ab}(\Phi_i) F_{kj}^b(\Phi_i) = 0 \Rightarrow \frac{d^2 f}{dr^2} + \frac{f(f^2 - 1)}{r^2} = 0. \quad (\text{B.6})$$

We can see three solutions of this equation  $f_1 = 0$ ,  $f_2 = +1$ ,  $f_3 = -1$  ( $r \neq 0$ ). The first solution corresponds to the naive unstable perturbation theory. Two nontrivial solutions are the Wu-Yang

monopoles [57] applied to the construction of physical variables in [58]. The Wu-Yang monopole is a solution of the classical equations everywhere besides the origin of the coordinates  $r = 0$ . The corresponding magnetic field is  $B_i^a(\Phi_k^{(0)}) = x^a x^i / (gr^4)$ . Following Wu and Yang [57], we consider the whole finite space volume, excluding an  $\epsilon$ -region around the singular point. To remove a singularity at the origin of coordinates and regularize its energy, the Wu-Yang monopole is considered as an infinite volume limit ( $V_0 \rightarrow \infty$ ) of the Bogomol'nyi-Prasad-Sommerfield (BPS) monopole [59]

$$f^{WY} = 1 \implies f^{BPS} = \left[ 1 - \frac{r}{\epsilon \sinh(r/\epsilon)} \right] \quad (\text{B.7})$$

with the finite energy  $\int d^3x [B_i^a(\Phi_k)]^2 \equiv V_0 < B^2 > = 4\pi/(g^2\epsilon) \equiv 1/(\alpha_s\epsilon)$  determining the finite volume regularization of the monopole size  $\epsilon = [\alpha_s V_0 < B^2 >]^{-1}$ . The vacuum energy-density of the monopole solution  $< B^2 >$  is removed by a finite counter-term in the Lagrangian.

The color particle excitations  $\mathcal{A}_\mu^{a(\text{rad})}$

$$\hat{A}_0(x_0, \vec{x}) = \hat{\mathcal{A}}_0^{(\text{rad})}(x_0, \vec{x}), \quad \hat{A}_k(x_0, \vec{x}) = \hat{\Phi}_k(\vec{x}) + \hat{\mathcal{A}}_k^{(\text{rad})}(x_0, \vec{x}). \quad (\text{B.8})$$

are considered as weak deviations in the vacuum background (B.5) by analogy with the Dirac radiation variables in QED (13), (14) with the constraint  $D_k^{ab}(\Phi_i) \mathcal{A}_k^{b(\text{rad})}(A) \equiv 0$  and the Gauss law

$$\Delta^{ab}(\Phi) \mathcal{A}_0^{b(\text{rad})} = -j_{t0}^{a(\text{rad})}, \quad (\text{B.9})$$

where  $\Delta^{ab}(\Phi) = [D_i^2(\Phi_k)]^{ab}$  is the Laplacian in the vacuum background, and  $j_{t0}^{a(\text{rad})}$  is the total color current.

The action can be expressed in terms of the radiation - type variables [21]

$$\begin{aligned} S_{\text{QCD}}|_{\frac{\delta S_{\text{QCD}}}{\delta A_0}=0} &\equiv S_{\text{QCD}}^{(\text{rad})} = \int d^4x \left[ \frac{1}{2} A_k^{a(\text{rad})} G_{ki}^{ab-1} A_i^{b(\text{rad})} + A_k^{(\text{rad})} j_k^{(\text{rad})} \right] \\ &+ (\psi^{(\text{rad})} \bar{\psi}^{(\text{rad})}, G_m) - \frac{1}{2} \int dx^0 d^3x d^3y j_{t0}^{a(\text{rad})}(x) V^{ab}(x-y) j_{t0}^{b(\text{rad})}(y) + \dots + S_{\text{soc.}}(\eta^{(\text{rad})}, \bar{\eta}^{(\text{rad})}), \end{aligned} \quad (\text{B.10})$$

where

$$S_{\text{soc.}}(\eta^{(\text{rad})}, \bar{\eta}^{(\text{rad})}) = \int d^4x \left[ \bar{\eta}^{(\text{rad})} \psi^{(\text{rad})} + \bar{\psi}^{(\text{rad})} \eta^{(\text{rad})} \right] \quad (\text{B.11})$$

is the source action,  $G_{ki}^{ab}$  and  $G_m$  are the Green functions of the gluon and quark in the background field  $\Phi_k^a$ ,

$$V^{ab}(\vec{z}) = -\frac{z^a(z)z^b(z)}{|\vec{z}|^2} \frac{1}{4\pi|\vec{z}|} + \sum_{\alpha=1,2} e_\alpha^a(z) e_\alpha^b(z) \left( d_- |\vec{z}|^{l_-} + d_+ |\vec{z}|^{l_+} \right)$$

is the potential [21, 60] considered as the Green function

$$[\Delta^{ab}(\Phi(\vec{z}))] V^{bc}(\vec{z}) = \delta^{ac} \delta^3(\vec{z}); \quad (\text{B.12})$$

here  $\vec{z} = \vec{x} - \vec{y}$ ,  $e_\alpha^a(z)$  are a set of orthogonal vectors in color space  $z_a e_\alpha^a(z) = 0$ ,  $d_\pm$  are constants, and  $l_+$ ,  $l_-$  can be found as roots of the “gold section” equation  $l^2 + l = 1$ :

$$l_+ = -\frac{1+\sqrt{5}}{2} \approx -1.618; \quad l_- = \frac{-1+\sqrt{5}}{2} \approx 0.618. \quad (\text{B.13})$$

The solution of the set of Schwinger – Dyson and Bethe – Salpeter equations for the rising potential was considered in numerous papers, where the effect of spontaneous chiral symmetry breaking was



described (see [20, 21, 61]). Therefore, this Green function contains the rising potential as the origin of the 'hadronization' of quarks and gluons in QCD [61].

The Wigner – Markov – Yukawa relativistic generalization of the instantaneous interaction in this case can lead to the bilocal theory with the spontaneous chiral symmetry breaking in the low-energy limit with the effective chiral Lagrangian one [34, 35].

## B.2. Zero mode of Gauss' constraint

The BPS - type regularization (B.7) of the monopole vacuum is interesting by the existence of a nonzero solution of homogeneous equation

$$[\Delta^{ab}(\Phi)]\mathcal{Z}^a = 0 \quad (\text{B.14})$$

of the type of the BPS solution [59]

$$\hat{\mathcal{Z}}(\vec{x}) = -i\pi \frac{\tau^a x^a}{r} f_0^{BPS}(r), \quad f_0^{BPS}(r) = \left[ \frac{1}{\tanh(r/\epsilon)} - \frac{\epsilon}{r} \right]. \quad (\text{B.15})$$

The complete solution of the Gauss equation (B.9), in this case, is the sum of a solution of the homogeneous equation (B.14) and a solution of the nonhomogeneous equation:

$$\mathcal{A}_0^{a(\text{rad})}(\vec{x}) = c(x_0)\mathcal{Z}^a - \int d^3y V^{bc}(\vec{x} - \vec{y}) j_0^{b(\text{rad})}(\vec{y}), \quad (\text{B.16})$$

where  $c(x_0)$  is the zero mode variable.

The BPS vacuum model gives the possibility to compare the radiation variables in the action (B.10) with the set of the Dirac - type variables removing the field  $A_0$  by the gauge transformations

$$\begin{aligned} \Phi_k^{(\text{D})} &= U^{(\text{D})}[\Phi_k^{(\text{rad})} + \partial_k]U^{(\text{D})-1}, \\ \mathcal{A}_l^{(\text{D})} &= U^{(\text{D})}[\mathcal{A}_l^{(\text{rad})}]U^{(\text{D})-1}, \\ \psi^{(\text{D})} &= U^{(\text{D})}\psi^{(\text{rad})}, \end{aligned} \quad (\text{B.17})$$

where

$$U^{(\text{D})} = \exp \left\{ \int^{x^0} d\bar{x}^0 c(\bar{x}^0) \hat{\mathcal{Z}}(\vec{x}) \right\} \equiv \exp \{ N(x^0) \hat{\mathcal{Z}}(\vec{x}) \}. \quad (\text{B.18})$$

The Dirac variables (B.17) (*dressed* by the zero mode phase factor) differ from the *radiation* ones in the action (B.10). The differences are Abelian U(1) anomaly and the new *gauge of physical sources*

$$S_{\text{soc.}}(\eta^{(\text{D})}, \bar{\eta}^{(\text{D})}) = \int d^4x \left[ \bar{\eta}^{(\text{D})} \psi^{(\text{D})} + \bar{\psi}^{(\text{D})} \eta^{(\text{D})} \right] \quad (\text{B.19})$$

as consequences of the gauge transformations (B.17).

The zero mode of the Gauss constraint (B.14) leads to the vacuum electric tension [62, 63]

$$F_{0i\text{vac}}^a = c(x_0) D_i^{ab}(\Phi) \mathcal{Z}^b \equiv \dot{N} D_i^{ab}(\Phi) \mathcal{Z}^b \quad (\dot{N} = \partial_0 N). \quad (\text{B.20})$$

The BPS monopole electric tension is proportional to the vacuum magnetic one. If we choose the coefficient of the proportionality as

$$D_i^{ab}(\Phi) \mathcal{Z}^b = B_i^a(\Phi) \frac{\alpha_s^2}{(2\pi)^2} \frac{1}{V_0 < B^2 >} \quad \left( \frac{g^2}{4\pi} \equiv \alpha_s \right), \quad (\text{B.21})$$

the zero mode  $c(x_0) = \dot{N}$  can be identified with a variable given by the winding number functional

$$\nu = \frac{g^2}{16\pi^2} \int_{x_{in}^0}^{x_{out}^0} dx^0 \int d^3x F_{\mu\nu}^a \tilde{F}^{a\mu\nu} = \frac{\alpha_s}{2\pi} \int d^3x D_i^{ab}(\Phi) \mathcal{Z}^b B_i^b(\Phi) \int_{x_{in}^0}^{x_{out}^0} dx^0 \dot{N} = N(x_{out}^0) - N(x_{in}^0)$$

showing that  $N$  can interpolate between different classical vacua like an instanton [51]<sup>11</sup>

$$N(x_{out}^0) = n_{out}, \quad N(x_{in}^0) = n_{in}. \quad (\text{B.22})$$

The equations (B.20) and (B.21) lead to the vacuum action of the zero mode dynamics

$$W_{\text{QCD}} = \int_{V_0} dx^0 \int d^3x \frac{1}{2} (F_{0i}^a)^2 = \int dx^0 \frac{\dot{N}^2}{2} \left( \frac{2\pi}{\alpha_s} \right)^2 \frac{1}{V_0 \langle B^2 \rangle}. \quad (\text{B.23})$$

As we pointed out above, one of consequences of the *dressed* variables (B.17) is the appearance of the Abelian anomaly [49] in the effective meson action obtained by the bosonization of QCD of the type of (A.11)

$$W_{eff} = \int dx^0 \left\{ \frac{1}{2} (\dot{\eta}_0^2 - m_0^2 \eta_0^2) V_0 + C_\eta \eta_0 \dot{N} \right\} \quad (\text{B.24})$$

in the pseudoscalar  $\eta_0$  meson isosinglet channel, where  $C_\eta = \frac{N_f}{F_\pi} \sqrt{\frac{2}{\pi}}$ ,  $N_f = 3$  is the number of flavors, and  $F_\pi = 92$  MeV is the weak decay coupling. This action is known as the Veneziano one [64].

The diagonalization of the total Lagrangian supplemented by the action (B.23) leads to an additional mass of the pseudoscalar  $\eta_0$  meson  $\Delta m_\eta^2 = \frac{N_f^2 \alpha_s^2 \langle B^2 \rangle}{F_\pi^2 2\pi^2}$  [21]. This result allows us to estimate the value of the vacuum chromomagnetic field in QCD [65]

$$\langle B^2 \rangle = \frac{2\pi^3 F_\pi^2 \Delta m_\eta^2}{N_f^2 \alpha_s^2} = \frac{0.06 \text{ GeV}^4}{\alpha_s^2}.$$

After calculation we can remove infrared regularization  $V_0 \rightarrow \infty$ .

Thus, there are observational arguments in favor of that physical fields are identified with the *dressed* one (B.17). However, the gauge of *dressed* sources (B.19) differ from the *radiation* ones (B.11) by the phase factors (B.18) depending on the winding number variable  $N$ . The initial position of winding number variable  $N$  are degenerated like the U(1) initial data  $e^{ix(\text{in})} = 1, x(\text{in}) = 2\pi k$ , where  $k$  is an integer number. The phase factors (B.18) are determined within the matrixes  $v^{(n)}(\vec{x}) = e^{n\hat{Z}(\vec{x})}$ , where  $n$  is an integer number which counts how many times a three-dimensional path  $v^{(n)}(\vec{x})$  turns around the  $SU(3)$ -manifold when the coordinate  $x_i$  runs over the space where it is defined. Even, if we choose the zero electric monopole energy fixing the zero momentum  $P_N = 0, N(\text{in}) = 0$ , we do not know a position of winding number variable due to this topological degeneration of all physical states with respect to the phase factor (B.17)  $U^{(D)} = e^{n\hat{Z}(\vec{x})}, n = 0, \pm 1, \pm 2, \dots$ . Therefore, the creation amplitude of the *dressed* color particles  $\psi^{(D)} = \psi^{(\text{rad})} U^D$  should be averaged over all degenerated states

$$\frac{1}{2L} \sum_{n=-L}^{n=+L} \langle \text{vacuum} | \psi^{(\text{rad})} U^D | q \rangle = \frac{e^{i\omega x_{in}^0}}{(2\pi)^{3/2} \sqrt{2\omega}} \frac{1}{2L} \sum_{n=-L}^{n=+L} e^{n\hat{Z}(\vec{x})} = 0, \quad (\text{B.25})$$

---

<sup>11</sup>In this case, the monopole interpolation amplitude  $\exp\{iP_N[N(x_{out}^0) - N(x_{in}^0)]\}$  coincides with the instanton one  $\exp\{-\frac{8\pi^2}{g^2}(n_{out} - n_{in})\}$  for a nonphysical value of the momentum  $P_N = i\frac{8\pi^2}{g^2}$ .

where  $|q\rangle$  is the Dirac one-particle state (B.17) with a quantum number  $q$ . Averaging over all parameters of the topological initial data can lead to a complete destructive interference of all color amplitudes [62] and the zero probability amplitude of creation of color quarks or gluons; whereas their *radiation* propagators have poles like the propagators of electron and photons and the colorless amplitudes of the type of the current production  $\langle j_\mu^{Da} j_\nu^{Da} \rangle = \langle j_\mu^{(\text{rad})a} j_\nu^{(\text{rad})a} \rangle$  do not disappear. In this case, only colorless (“hadron”) states have to form a complete set of physical states.

Disappearance of color physical states due to the topological degeneration (B.25) can be considered as color confinement, compatible with the quark – hadron duality as the accepted method of analysis of observational data [24].

## References

- [1] P.A.M. Dirac, Proc. Roy. Soc., **A 114**, 243–249 (1927);  
P.A.M. Dirac, Can. J. Phys., **33**, 650–661 (1955).
- [2] I.V. Polubarinov, Physics of Particles and Nuclei, **34**, 377–410 (2003).
- [3] L.D. Faddeev, Teor. Mat. Fiz. **1**, 3 (1969).
- [4] Nguyen Suan Han and V. N. Pervushin, Mod.Phys.Lett. **A2**, 367 (1987).
- [5] L. Faddeev and V. Popov, Phys. Lett. **B 25**, 29 (1967).
- [6] D. Bardin and G. Passarino, “The standard model in the making: precision study of the electroweak interactions”, Clarendon, Oxford, 1999.
- [7] F. Fock, Z. Phys., **39**, 226 (1926); *ibid.* , **57**, 261 (1929).
- [8] H. Weyl, Z. Phys., **56**, 330 (1929).
- [9] E.P. Wigner, Ann. of Math. **40**, 149 (1939).
- [10] S. Schweber, An Introduction to Relativistic Quantum Field Theory, Row, Peterson and Co, Evanston, Ill., Elmsford, N.Y, 1961.
- [11] N. N. Bogoliubov, A. A. Logunov, and I. T. Todorov, General Principles of Quantum Field Theory, Nauka, Moscow, 1969.
- [12] S.A. Gogilidze, V.N. Pervushin, and A.M. Khvedelidze, Physics of Particle and Nuclei, **30**, 66 (1999).
- [13] W. Kummer, Nuovo Cimento. **31**, 219 (1964); G. C. Wick, Phys. Rev. **96**, 1124 (1954);  
R. E. Cutkosky, Phys. Rev. **96**, 1135 (1954).
- [14] N. Nakanishi, Suppl. Progr. Theor. Phys. **43**, 43 (1969).
- [15] B. Zumino, J. Math. Phys., **1**, 1 (1960).
- [16] J. Schwinger, Phys. Rev., **127**, 324(1962).
- [17] L.D. Landau, E.M. Lifschitz, Theoretical Physics. Statistical Physics II. Moscow: Nauka, 1978.

- [18] M. Gell-Mann, K. Brueckner, Phys.Rev., **106**, 364 (1957).
- [19] L. Münchow, R. Reif, Recent Development in the Many Body Problem. Leipzig: Teubner-  
Texte Vol.7, 1985.
- [20] Yu. L. Kalinovsky et al., Sov. J. Nucl. Phys., **49**, 1709 (1989);  
V.N. Pervushin, Nucl. Phys. B (Proc. Supp.), **15**, 197–206 (1990);  
D. Blaschke et al., Nucl. Phys. A, **586**, 711 (1995).
- [21] V. N. Pervushin, Physics of Particles and Nuclei, **34**, 348 (2003); [hep-th/0109218];  
L.D. Lantsman and V.N. Pervushin, Physics of Atomic Nuclei, **66**, 1384 (2003).
- [22] Yu.L. Kalinovsky, L. Kaschluhn, V.N. Pervushin, Phys. Lett., 231 B, 288 (1989).
- [23] S. Love, Ann. Phys., **113**, 153 (1978).
- [24] R.P. Feynman, Photon Hadron Interaction, New-York, N.Y., 1972.
- [25] H.-P. Pavel and V.N. Pervushin, Int. J. Mod. Phys. **A 14**, 2885 (1999).
- [26] A.Z. Dubničková, S. Dubnička, V.N. Pervushin, and M. Sečanský, JINR, E2-2006-80,  
Dubna, 2006; hep-ph/0606005.
- [27] I. Mikulec, Kaon Physics, XXIV Physics in Collision - Boston, June 27-29, 2004.
- [28] G. Altarelli and L. Maiani, Phys. Lett. **52B**, 351 (1974).  
M. K. Gaillard and B. W. Lee, Phys. Rev. Lett. **33**, 108 (1974).
- [29] A. I. Vainshtein et al., Yad. Fiz. **24**, 820 (1976);  
S. S. Gershtein and M. Yu. Khlopov, JETP Lett., **23**, 338 (1976).
- [30] A. A. Belkov, Yu. L. Kalinovsky, V. N. Pervushin, Description of  $K^+ \rightarrow \pi^+ e^+ e^-$  Decay  
in Quantum Chiral Theory. JINR-P2-85-107, Feb. 1985. (Dubna, JINR);  
A. A. Bel'kov et al., Sov. J. Nucl. Phys. **44**, 690 (1986).
- [31] G. Ecker et al., Nucl. Phys. **B 291**, 692 (1987).
- [32] A. A. Bel'kov et al., Phys. Part. Nucl. **26**, 239 (1995).
- [33] A. A. Bel'kov et al., Phys. Lett. **220 B**, 459 (1989).
- [34] M. K. Volkov and V. N. Pervushin, Usp. Fiz. Nauk **120**, 363 (1976);  
M. K. Volkov and V. N. Pervushin, “Essentially Nonlinear Field Theory, Dynamical Sym-  
metry and Pion Physics”, Atomizdat, Moscow, ed. D. I. Blokhintsev, 1979 (in Russian).
- [35] J. Gasser and H. Leutwyler, Ann. of Phys. **158**, 142 (1984).
- [36] M. Volkov, V. Pervushin, Phys. Lett. **51 B**, 356 (1974).
- [37] Review of Particle Physics, Phys. Lett., **592 B**, 1 (2004).
- [38] M. E. Biagini, S. Dubnička, E. Etim, P. Kolar, Nuovo Cim., **A 104**, 363 (1991);  
S. Dubnička, A. Z. Dubničková, A. Antušek, Acta Physica Slovaca, **49**, 165 (1999);  
S. Dubnička, A. Z. Dubničková, J. Phys., **G28**, 2137 (2002).

- [39] V. N. Pervushin, Theor. Math. Phys., **27**, 330 (1977).
- [40] Yu. L. Kalinovsky, V. N. Pervushin, Sov. J. Nucl. Phys., **29**, 475 (1979).
- [41] P. Bloch et al., Phys. Lett., **56 B**, 201 (1975).
- [42] V.N.Pervushin, H. Reinhardt, D. Ebert, Sov. J. Part. Nucl., **10**, 1114 (1979).
- [43] M. A. Markov, J. Phys. (USSR), **2**, 453 (1940); H. Yukawa, Phys. Rev., **77**, 219 (1950).
- [44] J. Lukierski, M. Oziwicz, Phys. Lett., **69**, 339 (1977).
- [45] Yu.L. Kalinovsky, L. Kaschluhn, V.N. Pervushin, Phys.Lett. **B231**, 288 (1989).
- [46] A.A. Bogolubskaya, Yu.L. Kalinovsky, W. Kallies , V.N. Pervushin, Acta Phys. Polonica, **B 21**, 139 (1990).
- [47] Yu.L. Kalinovsky et al., Few Body System, **10**, 87 (1991).
- [48] M.K. Volkov Ann. Phys., **157**, 285 (1984).
- [49] D. Ebert and H. Reinhardt, Nucl. Phys., **B 271**, 188 (1986).
- [50] H. Fritzsch, M. Gell-Mann, and H. Leutwyler, Phys.Lett., **B 69**, 375 (1973).
- [51] A. A. Belavin et al. – Phys. Lett., **59**, 85 (1975);  
R. Jackiw, C. Rebbi, Phys. Lett., **63B**, 172 (1976);  
R. Callan, R. Dashen, D. Gross, Phys. Rev., **D 17**, 2717 (1977);
- [52] L. D. Faddeev, Proc. Fourt. Intern. Symposium on Nonlocal Field Theory (1976), D1-9788, Dubna, 1977, p.267.
- [53] R. Jackiw, Rev. Mod. Phys., **49**, 681 (1977).
- [54] G.'t Hooft, Phys. Rep., **142**, 357 (1986);  
G.'t Hooft, Monopoles, Instantons and Confinement, hep-th/0010225 (2000).
- [55] A.A. Vladimirov, D.V. Shirkov, Soviet. Phys. Usp., **22**, 860 (1979).
- [56] S.G. Matinyan, G K. Savidy, Nucl. Phys., **B 134**, 539 (1978).
- [57] T.T. Wu and C.N. Yang, Phys. Rev., **v.D 12** 3845 (1975).
- [58] L.D. Faddeev, and A.J. Niemi, Nature, **387** 58 (1997).
- [59] M.K. Prasad, C.M. Sommerfield, Phys. Rev. Lett., **35**, 760 (1975);  
E.B. Bogomol'nyi, Yad. Fiz, **24**, 449 (1976).
- [60] D. Blaschke, V. Pervushin, Röpke G., Proc. of Int. Seminar *Physical variables in Gauge Theories* Dubna, September 21-25, 1999 (E2-2000-172, Dubna, 2000 Edited by A. Khvedelidze, M. Lavelle, D. McMullan, and V. Pervushin) p.49; hep-th/0006249.
- [61] I.V. Puzynin et al., Physics of Particles and Nuclei, **30** 87 (1999).

- [62] V. N. Pervushin, Riv. Nuovo Cim., **8**, N 10, 1 (1985).
- [63] S. Gogilidze, Nevena Ilieva, V.N. Pervushin, Int. J. Mod. Phys. **A 14**, 3531 (1999).
- [64] G. Veneziano, Nucl. Phys., **B 195**, 213 (1979).
- [65] D. Blaschke, et al., Phys. Lett., **397 B**, 129 (1997).